



A Case of Incidentally Detected Gastric Outlet Obstruction in a Patient with Foreign Body Ingestion

**Anil Dev ^{a++*}, Ajitha M.B ^{b#}, Malla Balaji ^{ct}, Vijay A ^{dt},
Kesireddy Pavan ^{et} and Harshitha ^{ft}**

^a Department of General Surgery, Bangalore Medical College And Research Institute, 77-188/A1, APS Police Colony, Kallur Estates, Kallur, Kurnool-2, 518003, India.

^b Department of General Surgery, BMCRI, Address- #275,3rd B phase, UAS Layout, Shivanahalli, Yelahanka, Bengaluru-560064, India.

^c Department of General Surgery, Room no. 106, Sharavathi Boys Hostel, Bangalore Medical College and Research Institute, K.R. Market, Bangalore, Karnataka, 560002, India.

^d Department of General Surgery, Room no. 204, Sharavathi Boys Hostel, Bangalore Medical College and Research Institute, K.R. Market, Bangalore, Karnataka, 560002, India.

^e Department of General Surgery, Room no. 006, Sharavathi Boys Hostel, Bangalore Medical College and Research Institute, K.R. Market, Bangalore, Karnataka, 560002, India.

^f Department of General Surgery, Bangalore Medical College And Research Institute, door no 492,SS3 Sai mahima Residency,16th cross, Ideal Homes Town, RR Nagar, Bangalore, 560098, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/125944>

Case Report

Received: 29/08/2024

Accepted: 31/10/2024

Published: 11/11/2024

⁺⁺ S/O Deva Balaswamy, Post Graduate;

[#] HOD & Professor;

[†] Post Graduate;

^{*}Corresponding author: E-mail: anildeva789@gmail.com;

Cite as: Dev, Anil, Ajitha M.B, Malla Balaji, Vijay A, Kesireddy Pavan, and Harshitha. 2024. "A Case of Incidentally Detected Gastric Outlet Obstruction in a Patient With Foreign Body Ingestion". Asian Journal of Case Reports in Surgery 7 (2):511-16. <https://journalajcrs.com/index.php/AJCRS/article/view/576>.

ABSTRACT

Background: This case report discusses our experience of managing a case of metal key ingestion with incidentally detected gastric outlet obstruction by trial of endoscopy followed by surgery.

Presentation of Case: A case of 21 years old male prisoner presented with 2 days history of ingestion of metallic key and postprandial vomiting with history of mild abdominal distension and upper abdominal pain with no previous significant history. Clinical examination was unremarkable. Patient was kept nil per orally and serial Xray were taken which showed movement of foreign body following which CT abdomen and pelvis was done due to doubtful location in Xray. In CT over distended stomach measuring approximately 25 x 10 cm with metallic foreign body 4x2 cm noted in the stomach following which patient was posted for endoscopic retrieval of foreign body. In endoscopy, food stasis was noted in stomach with pinpoint pylorus, scope couldn't be negotiated beyond pylorus, key couldn't be visualised. Due to repeated vomiting and abdominal pain, patient was planned for emergency operation and underwent gastrotomy with key retrieval with gastrojejunostomy with truncal vagotomy with feeding jejunostomy. Patient had an uneventful recovery and was discharged on post operative day 25.

Discussion: Late-onset pyloric stenosis is due to the persistence of the infantile form, which becomes clinically evident only at a later stage, when a triggering event, like foreign body ingestion resulting inflammation, edema, or spasms, which precipitates pyloric occlusion [1]. Acute inflammation of pylorus can lead to gastric outlet obstruction manifested by early satiety, anorexia, weight loss, nausea, vomiting. In chronic inflammation, stomach can become massively dilated and lose its muscular tone. 12% of patients with peptic ulcer presented with gastric outlet obstruction as a direct consequence of a pyloric canal ulcer with associated pylorospasm [2].

Conclusion: Asymptomatic presentation of gastric outlet obstruction is rare, and hence, should be kept in mind in such situations of retained foreign body where it can cause acute inflammation and exacerbate gastric outlet obstruction.

Keywords: Key; endoscopy; gastric outlet obstruction; gastrojejunostomy.

1. INTRODUCTION

Foreign body ingestion in adults is relatively uncommon accounting to almost 15-20% of cases [3]. Unintentional ingestions are usually those of meat or fish bones. Intentional ingestion is mostly observed in people who are intellectually impaired, have psychiatric illness, prisoners or edentulous. Gastric outlet obstruction can be due to extrinsic and intrinsic factors. Rarely, a retained foreign body can cause acute inflammation and exacerbate gastric outlet obstruction [4]. According to American Society for Gastrointestinal Endoscopy guideline for management of ingested foreign bodies, 80%-90% of foreign bodies reaching the gastrointestinal tract will pass spontaneously, 10%-20% require endoscopic intervention and 1% require a surgical procedure [5,6]. We present you a case of foreign body ingestion with incidental finding of gastric outlet obstruction in a prison inmate.

2. PRESENTATION OF CASE

History: A 21 years old male prisoner with a history of two days old intentional ingestion of

metallic key presented to emergency department with complaints of multiple episodes of postprandial vomiting containing undigested food particles, with history of mild abdominal distension.

Clinical examination: soft, non-tender, mild distension, normal bowel sounds heard.

Investigations: Radiological investigations were done to confirm the foreign body. Serial radiographs were taken for first 24 hrs at an interval of 6hrs. Due to doubtful location on Xray, a CT scan was done.

In CT over distended stomach measuring approximately 25 x 10 cm with metallic foreign body 4x2 cm noted in the stomach followed by which patient was posted for esophagoduodenoscopy.

Management: A decision was made to remove the foreign body via endoscopic method under sedation. Food stasis was noted in stomach with pinpoint pylorus, and irregular inflamed surface in pylorus from where biopsy was sent. Key



Fig. 1. Serial erect X Ray abdomen showing foreign body movement



Fig. 2. CT abdomen and pelvis showing distended stomach with key in stomach



Fig. 3 and 4. Image showing retrieved stenosed area in pyloric region, key post retrieval



Fig. 5 and 6. Image showing gastrojejunostomy and feeding jejunostomy

couldn't be visualised and was suspected to be buried under inflamed mucosa. Scope couldn't be negotiated beyond pylorus.

Post endoscopy, patient was initially planned for second attempt of endoscopic retrieval but due to acute deterioration of patient with severe upper abdominal pain, with increasing abdominal distension and persisting vomiting and considering medicolegal implication of prisoner, patient was posted for emergency laparotomy where gastrotomy with retrieval of key with gastrojejunostomy with truncal vagotomy with distal feeding jejunostomy.

Histopathology report from stenosed site showed benign inflammatory lesion with features of both acute and chronic inflammation, suggesting probably a chronic ulcer causing

partial pyloric stenosis was triggered by ingestion of metallic key. Recovery was uneventful with superficial surgical skin infection drain removal on post op day 7, feeding jejunostomy on post op day 21 and discharged on post op day 25 after settlement of SSI and removal of all drains and sutures as per request of police incharge.

3. DISCUSSION

Management of foreign body ingestion is influenced by the patient's age and clinical condition, size, shape and classification of the ingested material, the anatomic location in which the object is lodged and the technical abilities of the endoscopist [7]. There are 3 sites where an object may fail to pass once it has negotiated cricopharynx, the upper oesophageal sphincter is

the narrowest part of the upper GI tract, the pylorus in the stomach, the ileocecal valve [8].

The persistence of the infantile form causes late-onset pyloric stenosis, which only manifests clinically later when a triggering event—such as ingesting a foreign body—causes inflammation, edema, or spasms that lead to pyloric occlusion.

Acute inflammation of the pylorus can lead to gastric outlet obstruction manifested by early satiety, anorexia, weight loss, nausea, vomiting. In chronic inflammation stomach can become massively dilated and lose its muscular tone. 12% of patients with a peptic ulcer presented with gastric outlet obstruction as a direct consequence of a pyloric canal ulcer with associated pylorospasm [9].

4. CONCLUSION

Gastric outlet obstruction triggered by acute events like foreign body ingestion is rare presentation. Once identified through imaging studies, prompt investigation should take place with EGD. We report a case of foreign body ingestion with an incidentally detected gastric outlet obstruction that was successfully treated with endoscopic/surgical procedure.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

CONSENT AND ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

REFERENCES

1. Iacoviello O, Verriello G, Castellaneta S, Palladino S, Wong M, Mattioli G, Giordano P, Francavilla R, Cristofori F et al. Case report: Late-onset hypertrophic pyloric stenosis in a 3-year-old boy: It is never too late. *Front Pediatr.* 2022 Aug 16;10:949144.
2. Costa CS, Pratas N, Capote H, et al. Massive gastric dilation caused by gastric outlet obstruction in the setting of peptic ulcer disease—A case report, *International Journal of Surgery Case Reports.* 2020;70:64-67.
3. Dhuha N. Boumarah, Lujain S. Binkhamis, Mohammed AIDuhileb et al. foreign body ingestion: Is intervention always a necessity? *Annals of Medicine and Surgery.* 2022;84:104944.
4. Kirk AD, Bowers BA, Moylan JA, et al. Toothbrush swallowing. *Arch Surg.* 1988;123:382-4.
5. American Society for Gastrointestinal Endoscopy. Guideline for the management of ingested foreign bodies. *Gastrointestinal Endoscopy.* 1995;42:622-5.
6. Eisen GM, Baron TH, Dominitz JA, et al. Guideline for the management of ingested foreign bodies. *Gastrointestinal Endoscopy.* 2002;55:802–6.
7. G, Wyllie R, Kay M. Foreign body ingestion in infants and children: location, location, location. *Clin Pediatr (Phila).* 2002;41:633–40.
8. Conners GP, Mohseni M. Pediatric foreign body ingestion. In: *Stat Pearls Treasure Island (FL): StatPearls;* 2024 Jan.
9. Trivarna Dey, Manjunath BD, Niyaz Ahmed et al. A case report on challenges in retrieving an ingested mobile phone. *AJCRS* 120553, 2024;7(2):378-82.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/125944>