



Barriers and Determinants of Insecticide-treated Net Utilization for Malaria Prevention among Pregnant Women in Tamale Metropolis, Ghana

**Suraya M. Mohammed ^{a*}, Delilah Nonterah ^a,
Ruhaima Abukari ^b, Ummul-Kulsum Issahaku ^b,
Jelila Mumuni ^a, Suweiba Hamidu ^a, Fati Labaran ^c,
Habib Mahama ^d, Nixsion A. Niadawe ^e,
Karima Mohammed ^f, Ramatu Akunvane ^c,
Manzuk Ebenezer ^e, Seidu Balkisu ^a, Issaka Zuwera ^a,
Abukari Nashiru ^g and Alhassan Abdul-Rahaman ^g**

^a Ministry of Health, Nurses' and Midwives' Training College, P.O. Box 565, Tamale, Northern Region, Ghana.

^b Nurses' and Midwives' Training College, Kpembe, P. O. Box SL 98, Salaga, Savannah Region, Ghana.

^c Nursing and Midwifery Training College, Zuarungu, P. O. Box 660, Bolgatanga, Upper East Region, Ghana.

^d Ghana Health Service, Kassena Nankana Municipal Health Directorate, P. O Box 8, Navrongo, Upper East Region, Ghana.

^e Community Health Nurses Training College, Navrongo, P. O. Box 185, Navrongo, Upper East Region, Ghana.

^f Nursing and Midwifery Training College, Gushegu, P. O. Box GU 13, Gushegu, Northern Region, Ghana.

^g College of Nursing and Allied Health Sciences, P.O. Box 10, Nalerigu, North-East Region, Ghana.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/sajp/2024/v7i4205>

*Corresponding author: E-mail: smohammedmalle@nmtctamale.edu.gh;

Cite as: Mohammed, Suraya M., Delilah Nonterah, Ruhaima Abukari, Ummul-Kulsum Issahaku, Jelila Mumuni, Suweiba Hamidu, Fati Labaran, Habib Mahama, Nixsion A. Niadawe, Karima Mohammed, Ramatu Akunvane, Manzuk Ebenezer, Seidu Balkisu, Issaka Zuwera, Abukari Nashiru, and Alhassan Abdul-Rahaman. 2024. "Barriers and Determinants of Insecticide-Treated Net Utilization for Malaria Prevention Among Pregnant Women in Tamale Metropolis, Ghana". South Asian Journal of Parasitology 7 (4):411-26. <https://doi.org/10.9734/sajp/2024/v7i4205>.

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/127714>

Original Research Article

Received: 10/10/2024

Accepted: 12/12/2024

Published: 16/12/2024

ABSTRACT

Introduction: Insecticide-treated mosquito nets (ITNs) are proven malaria prevention tools, particularly for pregnant women who are at increased risk of malaria-related complications. In Ghana, malaria remains a leading cause of morbidity and mortality among pregnant women, making the use of ITNs critical in reducing malaria incidence. Aim: This study aimed to assess the barriers and determinants of ITNs utilization among pregnant women in the Tamale Metropolis. Methods: A cross-sectional survey was conducted with 399 pregnant women in the Tamale Metropolis using stratified sampling. A structured questionnaire solicited responses from the participants. Data were analyzed using Stata version 18, and a p-value less than 0.05 was considered statistically significant.

Results: The study found that 98.5% of participants were aware of ITNs, with the majority gaining knowledge through mass media (81.7%) and antenatal care centers (79.1%). Despite this high level of awareness, 92.5% of participants owned at least one ITN. The main source of ITNs was free house-to-house distribution (79.7%) and ANC centers (78.6%). However, while 82.7% of participants used ITNs at home, only 22.6% reported consistent nightly use. In terms of education level, basic education (AOR=0.06, 95% CI: 0.02-0.18), secondary education (AOR=0.08, 95% CI: 0.03-0.23), and tertiary education (AOR=0.29, 95% CI: 0.09-0.95, $p=0.042$) had lower odds compared to those with no formal education. Respondents who were self-employed (AOR=0.04, 95% CI: 0.01-0.20) and unemployed (AOR=0.01, 95% CI: 0.005-0.32, $p<0.001$) had reduced odds compared to those who were employed. Additionally, household monthly income was significantly associated with those earning 500-999 (AOR=0.07, 95% CI: 0.02-0.21, $p<0.001$), 1000-1999 (AOR=0.10, 95% CI: 0.01-0.30, $p<0.001$), and ≥ 2000 (AOR=0.10, 95% CI: 0.03-0.21, $p<0.001$) had lower odds compared to the lowest income category (0-499).

Conclusions: While ownership of ITNs is high, consistent use among pregnant women remains low due to practical barriers such as discomfort and lack of proper facilities. Addressing these issues through education, infrastructural improvements, and increased access to ITNs, especially in remote areas, could enhance ITN usage.

Keywords: *Insecticide-treated mosquito nets; malaria prevention; pregnant women; utilization; tamale metropolis; Ghana; barriers; socioeconomic factors.*

1. INTRODUCTION

In 2022, an estimated 249 million malaria cases and 608,000 malaria-related deaths were reported across 85 countries worldwide [1]. The WHO African Region bore the brunt of this burden, accounting for 94% (233 million) of global malaria cases and 95% (580,000) of malaria deaths. Alarming, children under the age of five represented approximately 80% of all malaria deaths in the Region, highlighting the severe impact of the disease on vulnerable populations [1]. Malaria is a primary contributor

to morbidity, mortality, poverty, and diminished productivity among the majority of individuals residing in low- to middle-income countries (LMICs) [2]. Malaria elevates the risk of severe pregnancy problems for the mother, fetus, and neonate. Insecticide-treated bed nets (ITNs) are a kind of personal protection that reduces malaria infection, severe illness, and mortality [3]. The efficient application of ITNs has diminished malaria morbidity by 50% in children under 5 years of age and decreased mortality by 17% [4,5]. The WHO Roll Back Malaria (RBM) initiative, initiated in 1998, aimed to prevent and

manage malaria in pregnant women through the utilization of insecticide-treated nets (ITNs) and additional strategies [6,7].

Using insecticide-treated nets (ITNs) is an intervention demonstrated to diminish malaria infections and decrease malaria-related mortality in endemic areas [8]. A systematic review of ITN use revealed that ITN use revealed that ITNs lessen the prevalence of *Plasmodium falciparum* malaria by 17% [9]. In Ghana, access to ITNs in households resulted in a 7.1 percentage point decrease in self-reported malaria among women [9]. Several reasons contribute to disparities in the utilization of ITNs. A study conducted in Ghana by Bawuah and Ampaw [10] demonstrated that dwelling location and socioeconomic status are predictors of ownership and utilization of insecticide-treated nets (ITNs). Luukwa et al. [11] contended that children and secondary-educated pregnant women from impoverished rural families in a specific region of Ghana exhibited the highest utilization of ITNs compared to pregnant women with differing attributes [12]. In Malawi, the likelihood of utilizing ITNs was low among uneducated women, those with primary education, and impoverished individuals [13].

Ghana, ranked among the fifteen nations with the highest malaria infection rates globally, experienced a 5% rise in absolute case numbers from 2017 to 2018, resulting in almost 0.5 million additional cases [8]. The nation exhibits the highest prevalence of Intermittent Preventive Treatment in Pregnancy (IPTp) among pregnant women at 78%, with the proportion of those receiving the third dosage rising from 39% in 2016 to 60% in 2018, and reaching 61% in 2019 [14]. IPTp entails administering a therapeutic dose of an efficacious antimalarial medication to all pregnant women without prior confirmation of malaria infection [15,16].

In Ghana, the deliberate promotion and intervention about ITNs have led to an increase in household ownership of ITNs from 49% in 2011 to 68% in 2014 [17,18]. Nonetheless, the scarcity of access and knowledge on ITNs and malaria has been recognized as a substantial obstacle to ITN adoption in Africa. Over 40% of insecticide-treated nets (ITNs) in households remain unutilized [19]. Certain pregnant women forgo the use of bed nets owing to financial limitations, lost opportunities during

complimentary distributions, discomfort from heat and odor, and challenges associated with suspending the net [12,17].

Notwithstanding advancements in preventive and therapeutic strategies, malaria continues to be widespread in Ghana, constituting 38.1% of all outpatient department cases and over half of all hospital admissions for children under five years of age. Ghana is striving for universal ITN coverage, necessitating ongoing monitoring and evaluation of access to and utilization of treated bed nets to inform malaria control policies and practices, especially in high-risk regions of the country. In the Tamale Metropolis of northern Ghana, where malaria transmission is prevalent, comprehending the determinants affecting malaria prevention and the utilization of insecticide-treated nets (ITNs) among pregnant women is essential for enhancing mother and child health outcomes. Inadequate use of ITNs, together with insufficient understanding or misconceptions regarding malaria prevention, significantly endangers pregnant women and their unborn children, potentially resulting in issues such as anemia, low birth weight, and even maternal and fetal mortality. This study seeks to evaluate barriers and determinants of ITNs knowledge, ownership, and utilization of ITNs among pregnant women in the Tamale Metropolis. The findings will elucidate deficiencies in existing interventions and assist in formulating more efficacious malaria preventive measures specifically designed for pregnant women in the Tamale Metropolis.

2. METHODOLOGY

2.1 Study Setting

The Tamale Metropolitan Assembly (TMA) is a part of the 261 Metropolitan, Municipal and District Assemblies (MMDAs) in Ghana, located in the Northern Region. Elevated to the status of Metropolis in 2004, it is the capital of the Metropolis and is situated between latitude 9.16° and 9.34° North and longitudes 00.36° and 00.57. The topography generally rolls with shallow valleys and isolated hills, but they do not hinder physical development. The Metropolis shares boundaries with Savelugu Municipality, Yendi Municipal Assembly, Tolon District, Central Gonja District, and East Gonja Municipal. The 2021 population and housing census recorded a population of 374,744.

2.2 Study Design

The study used cross-sectional study with an a cross-sectional study with an emphasis on quantitative research. Cross-sectional studies observe factors without altering them and collect data from numerous people at once. Cross-sectional study designs allow researchers to concurrently examine several characteristics, including age, gender, income, and education level, in connection to the utilization of insecticide-treated mosquito nets.

2.3 Study Population

The study population included pregnant women attending antenatal care (ANC) in the Tamale Metropolis.

2.4 Inclusion and Exclusion Criteria

2.4.1 Inclusion criteria

The inclusion criteria included the following:

1. Pregnant women residing within the Tamale Metropolis.
2. Women who are currently attending antenatal care (ANC) clinics or are reachable within health facilities in the Tamale Metropolis.
3. Pregnant women who provide informed consent to participate in the study.
4. Women aged 18 years and above.

2.4.2 Exclusion criteria

1. Non-pregnant women or women who have already given birth and are in the postpartum period.
2. Pregnant women who do not reside within the Tamale Metropolis.
3. Women with serious medical conditions or complications that prevent them from participating in the study.
4. Pregnant women who decline to give consent for participation.

2.5 Sample Size Determination

The formulae [20] was used to estimate the sample size.

$$n = \frac{z^2 \sigma}{m_0 E^2}$$

- Sample size (n) =?
- Variance (σ) = P (1-P)
- A recent student study in Ghana estimated the utilization of ITNs is 62.0% [21]. Thus, $p = 62\%$ (0.62)

Thus, Variance (σ) = P (1-P)

Variance (σ) = 0.62(1-0.62)

$\sigma = 0.2356$

- Margin of error (MOE)=0.05
- The critical value (Z) at 95% Confidence Interval = 1.96

$n = 363$

To account for noncompliance and response rate, 10.0% of the estimated was added. Thus, the sample size for this study was 399.

2.6 Sampling Techniques

A stratified sampling technique was used to select research participants from four healthcare institutions in the Tamale Metropolis. Stratified sampling was selected to guarantee that the sample correctly reflected pregnant women from each institution, facilitating proportional representation and augmenting the dependability of the results. The research population was stratified according to the four healthcare institutions. Participants were randomly chosen from each stratum according to the number of pregnant women receiving prenatal care at each institution. This strategy guaranteed the capture of diversity within each institution, providing a thorough knowledge of the determinants affecting postnatal care consumption in the Tamale Metropolis.

2.7 Data Collection Instruments

This study used A structured questionnaire as the primary data collection instrument. The questionnaire was designed to collect quantitative data and was developed by reviewing relevant literature and incorporating elements from existing validated instruments [8,22–24]. The questionnaire was structured according to the study objectives and included the following sections: Section A consisted of the socio-demographic characteristics, Section B included the knowledge of malaria prevention during pregnancy, Section C focused on accessibility and ownership of ITNs, Section D had questions on the utilization of the ITNs and Section E on the barriers to using the ITNs.

2.8 Data Collection Procedures

Data collection was executed by a team of five trained field enumerators. Before data collection, the enumerators received extensive training and

participated in role-playing exercises to guarantee uniformity and standardization in giving the questionnaire. This preparation sought to improve the dependability and precision of the data obtained.

Authorization to do the research was secured from each of the four designated healthcare institutions within the Tamale Metropolis. Moreover, informed permission was obtained from all participants before their involvement. The enumerators performed in-person interviews with the participants, each lasting between 20 to 30 minutes.

Confidentiality was rigorously maintained throughout the data gathering procedure, safeguarding all participants' information. Ethical concerns were maintained by following the principles of voluntary participation, informed consent, and safeguarding the respondents' privacy and rights.

2.9 Data Analysis and Presentation

Data cleaning was performed using Microsoft Excel to ensure accuracy and completeness before being exported to STATA version 18 for analysis. Both descriptive and inferential statistics were conducted to address the specific objectives of the study. The results were presented in tabular form for clarity.

The utilization of insecticide-treated nets (ITNs) was defined as respondents who reported consistent use of ITNs for malaria prevention. To

assess the association between ITN utilization and demographic factors, as well as health accessibility characteristics, chi-square tests were conducted. Statistical significance was determined at a threshold of $p < 0.05$.

3. RESULTS

3.1 Socio-demographic Characteristics

Most participants are aged 20-29 (45.1%) and have no education (33.8%). Over half have fewer than three children (54.9%) and follow Islam (69.9%). Most are married (65.9%), with most respondents being unemployed (43.6%). Regarding health, more than half (52.4%) experienced more than one malaria episode during their pregnancy. Household income levels showed that 34.3% earn less than 500 Ghana cedis monthly (Table 1).

3.2 Knowledge of Malaria Prevention During Pregnancy

Almost all participants (98.5%) were aware of insecticide-treated nets (ITNs), with the primary sources of this knowledge being mass media (81.7%) and antenatal care (ANC) centers (79.1%). Most participants recognized the importance of ITNs for reducing contact with mosquitoes (90.6%) and preventing malaria infection (81.4%). Other common preventive methods reported include using insect repellents (72.9%), avoiding stagnant water (74.7%), and covering exposed skin at night (74.9%) (Table 2).

Table 1. Socio-demographic characteristics

Variable	Category	Frequency	Percentage
Age	<20	43	10.8
	20-29	180	45.1
	30-39	129	32.3
	≥ 40	47	11.8
Highest level of education	No formal education	135	33.8
	basic education	83	20.8
	secondary	93	23.3
	tertiary	88	22.1
Parity	<3	219	54.9
	≥ 3	180	45.1
Religion	Islam	279	69.9
	Christian	113	28.3
	Traditional	7	1.8

Variable	Category	Frequency	Percentage
Marital status	Single	102	25.6
	Married	263	65.9
	Divorced	30	7.5
	widowed	4	1.0
Occupation	Unemployed	174	43.6
	self-employed	102	25.6
	employed	123	30.8
Malaria episodes during current pregnancy	once	190	47.6
	More than once	209	52.4
Household monthly income(Ghana cedis)	0-499	137	34.3
	500-999	129	32.3
	1000-1999	56	14.0
	≥ 2000	77	19.3

Table 2. Knowledge of malaria prevention during pregnancy

Variable	Category	Frequency	Percentage
Knowledge on ITNs	Yes	393	98.5
	No	6	1.5
Source of knowledge	ANC centers	311	79.1
	Health education campaigns	249	63.4
	Mass media	321	81.7
	Friends and colleagues	138	35.1
	Family	174	44.3
	Neighbors	38	9.7
Opinion on importance of ITNs during pregnancy	They reduce contact between pregnant women and mosquitoes	356	90.6
	They provide a shelter for mosquitoes	4	1.0
	They drive away mosquitoes from the room	94	23.9
	It is an instruction from the ANC clinic that must be followed	262	66.7
	They help prevent malaria infection	320	81.4
Other ways of preventing malaria in pregnancy	Use of insect repellents	291	72.9
	Indoor residual spraying	248	62.2
	Taking antimalaria drugs	47	11.8
	Avoiding stagnant water near the house	298	74.7
	Covering exposed skin at night	299	74.9
	Others(clearing bushes)	9	2.3

3.3 Accessibility and Ownership of ITNs

The study shows that 92.5% of participants own at least one ITN, with most obtaining them from free house-to-house distribution (79.7%) and ANC centers (78.6%). Among those who owned ITNs, ownership is relatively evenly distributed, with 32.8% owning one, 33.3% owning two, and

32.0% owning three or more. Most participants (76.2%) reported that ITNs were readily available when needed, though a substantial portion (39.6%) live 6-10 km from the nearest facility where they can obtain an ITN. Only 11% of participants expressed a specific preference for the type of ITN, with the most common preference being for a smooth texture (45.5%).

Table 3. Accessibility and ownership of ITNs

Variable	Category	Frequency	Percentage
Own at least one ITN	Yes	369	92.5
	No	30	7.5
Number of ITNs owned currently	1	121	32.8
	2	123	33.3
	3 or more	118	32.0
Source of ITN	ANC centers	290	78.6
	Market	33	8.9
	friends/relatives	35	9.5
	free house-house distribution	294	79.7
	pharmacy	45	12.2
ITNs were readily available when needed	Yes	304	76.2
	No	95	23.8
Distance to the nearest facility where ITNs can be obtained	<1 km	49	12.3
	1-5 km	125	31.3
	6-10 km	158	39.6
	>10 km	67	16.8
Have a preference of ITN	Yes	44	11.0
	No	355	89.0
ITN preferences	Smooth texture	20	45.5
	Reusable ITN	16	36.4
	Low chemicals ITN	8	18.2

3.4 Utilization of ITNs

The study found that 82.7% of participants used an ITN at home, only 22.6% reported consistent nightly use, and 60.6% used it the previous night. Many participants (43.9%) began using ITNs before pregnancy, while others started during pregnancy (37.3% in the first trimester). The

frequency of ITN use varies, with only 21.2% always using it. Most participants (70%) adequately placed the ITN by hanging it over the bed and tucking it under the mattress. However, consistent use is hindered by issues like discomfort due to heat (76.7%), inability to use it when it's hot (64.6%), and lack of proper facilities to hang the ITN (19.6%) (Table 4).

Table 4. Utilization of ITNs

Variable	Category	Frequency	Percentage
Utilize ITN at home.	Yes	330	82.7
	No	69	17.3
Consistently utilize ITN every Night	Yes	90	22.6
	No	309	77.4
Utilized ITN last night	Yes	200	60.6
	No	130	39.4
Started using ITNs	Before pregnancy	145	43.9
	During 1st trimester	123	37.3
	During 2nd trimester	26	7.9

Variable	Category	Frequency	Percentage
	During 3rd trimester	36	10.9
Frequency of ITN utilization			
	Always	70	21.2
	Often	69	20.9
	Sometimes	110	33.3
	Rarely	81	24.5
How do you usually place ITNs			
	Hanging it over the bed and tucking it under the mattress to prevent contact with the body	231	70.0
	Hanging it at one corner of the room	7	2.1
	Hanging it over the bed but the net has contact with the skin while sleeping under it	42	12.7
	Hanging over the windows and doors to prevent mosquitoes from entering	57	17.3
Sleep under ITNs even when it is hot.			
	Yes	122	37.0
	No	208	63.0
Reason for inconsistent use of ITNs			
	It is uncomfortable	113	47.1
	It causes excessive heat and discomfort	184	76.7
	I don't know how to hang it properly	42	17.5
	No facility to hang it in my room	47	19.6
	Can't use it when the weather is hot	155	64.6
	Traps mosquitoes inside when tucked in	4	1.7
	Heard it harms the baby in the womb	46	19.2
	Others(contains chemicals)	5	2.1

3.5 Barriers to Using ITNs

The primary barriers to using ITNs among pregnant women were excessive heat and discomfort (97.0%), with additional challenges such as difficulty using ITNs during hot weather (70.3%) and lack of knowledge on how to hang them properly (37.6%). Structural issues like the absence of facilities to hang ITNs (29.4%) and the discouraging routine of raising the net each morning (34.2%) also contributed to inconsistent use. Affordability was a concern for 20.9% of participants, and 26.7% reported not receiving an ITN when they were distributed for free. Suggested improvements to encourage ITN usage included making them readily available (80.6%), designing them for better air circulation (100.6%), providing adequate information on proper use (76.7%), and enhancing education on their benefits and safety for pregnant women (Table 5).

3.6 Association between Socio-demographic Characterize and Consistent Usage of ITNs

Several key variables showed significant associations in the study on consistent usage of insecticide-treated nets (ITNs). Age had a p-

value of less than 0.001, indicating younger individuals were less likely to use ITNs. Parity ($p = 0.012$) suggested that women with fewer than three children had higher usage rates. Educational level was highly significant ($p < 0.001$), with those holding tertiary degrees more likely to use ITNs. Marital status ($p = 0.012$) and occupation ($p = 0.004$) also correlated with ITN usage, as married and employed individuals reported higher usage. Furthermore, the number of malaria episodes ($p < 0.001$) and household income ($p = 0.013$) were significant, with multiple malaria episodes and lower-income households linked to more consistent ITN use (Table 6).

3.7 Determinant of ITNs among Pregnant Women

Several factors showed notable associations with the outcome. In terms of education level, individuals with basic education (AOR=0.06, 95% CI: 0.02-0.18, $p<0.001$), secondary education (AOR=0.08, 95% CI: 0.03-0.23, $p<0.001$), and tertiary education (AOR=0.29, 95% CI: 0.09-0.95, $p=0.042$) had lower odds compared to those with no formal education. Employment status also demonstrated significance, where self-employed individuals (AOR=0.04, 95% CI: 0.01-0.20, $p<0.001$) and unemployed individuals

(AOR=0.01, 95% CI: 0.005-0.32, $p<0.001$) had reduced odds compared to those who were employed. Additionally, household monthly income was significantly associated; those earning 500-999 (AOR=0.07, 95% CI: 0.02-0.21, $p<0.001$), 1000-1999 (AOR=0.10, 95% CI: 0.01-0.30, $p<0.001$), and ≥ 2000 (AOR=0.10, 95% CI: 0.03-0.21, $p<0.001$) had lower odds compared to the lowest income category (0-499) (Table 7).

Table 5. Barriers to using ITNs

Variable	Category	Frequency	Percentage
Challenges faced when using ITNs			
	Too costly, I cannot afford it	69	20.9
	Causes excessive heat and discomfort	320	97.0
	I don't know how to hang it properly	124	37.6
	No facility to hang it in my room	97	29.4
	I can't use it when the weather is hot	232	70.3
	The routine of raising it every morning after use is discouraging	113	34.2
	I heard it hurts the baby in the womb	78	23.6
	Did not get one when it was freely distributed	88	26.7
	Traps mosquitoes inside when tucked in	13	3.9
	Others(contains chemicals)	5	1.5
Measures to make ITN usage easy and correctly			
	Adequate information on how to hang ITNs should be provided	253	76.7
	They should be made readily available	266	80.6
	They should be given free of charge	235	71.2
	The price should be reduced	53	16.1
	Better design to allow air circulation	332	100.6
	Improved education on the benefits and safety of ITNs for pregnant women	228	69.1

Table 6. Association between socio-demographic characterize and consistent usage of ITNs

Variable	Category	Consistently usage of ITNs		P-value
		Yes	No	
Age				$P<0.001$
	<20	4(9.3%)	39(90.7%)	
	20-29	33(18.3%)	147(81.7%)	
	30-39	48(37.2%)	81(62.8%)	
	≥ 40	5(10.6%)	42(89.4%)	
Parity				$P=0.012$
	< 3	39(17.8%)	180(82.2%)	
	≥ 3	51(18.3%)	129(71.7%)	
Highest education level				$P<0.001$
	Tertiary	28(31.8%)	60(68.2%)	
	Secondary	11(11.8%)	82(88.2%)	
	Basic education	7(8.4%)	76(91.6%)	
	No formal education	44(32.6%)	91(67.4%)	
Religion				$P=0.260$
	Islam	67(24.0%)	212(76.0%)	
	Christianity	23(20.4%)	90(79.6%)	
	Others	0(0.0%)	7(100.0%)	
Marital status				$P=0.012$
	Married	64(24.3%)	199(75.7%)	
	Single	26(25.5%)	76(74.5%)	
	Divorced	0(0.0%)	30(100.0%)	
	Widowed	0(0.0%)	4(100.0%)	

Variable	Category	Consistently usage of ITNs		P-value
		Yes	No	
Occupation	Employed	28(22.8%)	95(77.2%)	P=0.004
	Self employed	34(33.3%)	68(66.7%)	
	Unemployed	28(16.1%)	146(83.9%)	
Malaria episodes during current pregnancy	Once	43(22.6%)	147(77.4%)	P<0.001
	More than once	47(22.5%)	162(77.5%)	
Aware of ITN	Yes	90(22.9%)	303(77.1%)	P=0.183
	No	0(0.0%)	6(100.0%)	
Household monthly income	0-499	42(30.7%)	95(69.3%)	P=0.013
	500-999	21(16.3%)	108(83.7%)	
	1000-1999	15(26.8%)	41(73.2%)	
	≥ 2000	12(15.6%)	65(84.4%)	

Table 7. Determinant of ITNs among pregnant women

Variable	Category	AOR (95% CI)	p-value
Age	<20	Ref*	
	20-29	0.58(0.14-2.38)	0.448
	30-39	2.93(0.49-17.65)	0.24
	≥ 40	0.31(0.04-2.61)	0.284
Highest level of education	No formal education	Ref*	
	Basic education	0.06(0.02-0.18)	p<0.001
	Secondary	0.08(0.03-0.23)	p<0.001
	Tertiary	0.29(0.09-0.95)	0.042
Marital status	Married	Ref*	
	Single/divorced/separated	1.32(0.59-2.94)	0.496
Parity	< 3	Ref*	
	≥ 3	2.16(0.65-7.22)	0.209
Occupation	Employed	Ref*	
	Self-employed	0.04(0.01-0.20)	p<0.001
	Unemployed	0.01(0.005-0.32)	p<0.001
Malaria episodes during current pregnancy	Once	Ref*	
	More than once	1.60(0.78-3.27)	0.199
Household monthly income	0-499	Ref*	
	500-999	0.07(0.02-0.21)	p<0.001
	1000-1999	0.10(0.01-0.30)	p<0.001
	≥ 2000	0.10(0.03-0.21)	p<0.001

Ref*= Reference, AOR-Adjusted Odds Ratio

4. DISCUSSION

The study assessed the utilization of ITNs among pregnant women in the Tamale Metropolis.

The study showed that majority of the respondents were aware of ITNs. Similar levels of ITN awareness have been reported in studies conducted in sub-Saharan Africa. For instance, a study by Asumah et al., [8] indicated that over

90% of households in regions targeted by malaria control programs were aware of ITNs. The widespread dissemination of malaria prevention messages via mass media and health centers supports such high awareness levels. The role of mass media and ANC clinics as key information sources is consistent with findings in the literature. A study by Saha et al. [25] demonstrated that radio campaigns and health facility visits during pregnancy significantly increased ITN awareness among pregnant women. ANC visits are particularly effective because they target women during a crucial stage when they are more likely to prioritize preventive health measures for themselves and their unborn children [26].

Recognition of ITNs for reducing mosquito contact and preventing malaria (90.6% and 81.4%, respectively) is comparable to findings in studies where ITNs are often considered the gold standard for malaria prevention [27–29]. The emphasis on the multiple advantages of ITNs corresponds with community-wide educational initiatives that present ITNs as both protective and economically viable. The application of repellents, the avoidance of stagnant water, and the covering of exposed skin at night conform to WHO-recommended malaria prevention strategies. Research has seen analogous tendencies, emphasizing the significance of multimodal preventative strategies, especially in regions where access to ITNs may be restricted or behavioral patterns affect compliance [30, 31].

While the findings broadly agree with existing research, there are nuanced areas of convergence and divergence. Near-universal awareness (98.5%) among participants is consistent with reports from regions with robust malaria control programs. This demonstrates the success of campaigns and ANC integration, similar to results from programs in Africa [32]. The prominence of ANC as a source reflects the increasing focus on maternal health as a critical entry point for malaria prevention efforts, as seen in programs by Roll Back Malaria (RBM). However, the literature often highlights gaps between ITN awareness and actual usage. Some studies indicate that factors such as affordability, accessibility, and cultural perceptions limit ITN use despite widespread awareness [33,34]. Despite participants' general comprehension of the advantages of ITNs, the marginally reduced awareness of malaria prevention (81.4%) relative to mosquito contact reduction (90.6%) may indicate insufficient messaging. Previous

research emphasizes comprehensive malaria education's significance in strengthening the connection between mosquito bites and malaria transmission [35]. Targeted campaigns, such as mass media and ANC programs, effectively reach vulnerable populations, particularly pregnant women and children. Recognizing ITNs and other preventive methods is influenced by culturally appropriate messaging and community engagement. However, socioeconomic factors, cultural norms, and healthcare access variations can explain differences in awareness and perceived importance of ITNs. Economic barriers, misconceptions about safety, and discomfort in hot climates can also contribute to discrepancies [36].

The study's finding that 92.5% of participants own at least one ITN is in line with other research that has found widespread ITN distribution efforts have been successful in increasing ownership. For example, research in sub-Saharan Africa has shown that ITN ownership rates often exceed 90% in areas with large-scale distribution programs [31]. The high rate of ITN ownership observed in this study is likely the result of effective interventions, such as free house-to-house distribution and ANC center distributions, which have been documented as effective strategies to increase access to ITNs [8]. However, the finding that 39.6% of participants live 6-10 km from the nearest ITN distribution facility highlights access barriers that may still exist, especially in rural or hard-to-reach areas. This is consistent with the literature, which suggests that even when ITNs are distributed freely, geographic barriers or poor infrastructure may hinder their accessibility for certain populations [8].

The study's finding that 82.7% of participants use ITNs at home but only 22.6% use them consistently every night is also consistent with previous studies, which have shown a gap between ITN ownership and consistent usage. For example, researches [8,37] found that while a high percentage of households owned ITNs, consistent nightly use often fell short, primarily due to discomfort, heat, and inadequate knowledge on proper usage. The study's finding that 76.7% of participants reported discomfort due to heat, and 64.6% had difficulty using ITNs when it was hot, supports these reports, suggesting that environmental factors significantly impact ITN use, especially in hot climates. Studies have consistently pointed out that discomfort during warm weather is a primary

barrier to ITN use, particularly in tropical areas where temperatures can make sleeping under a net challenging [38–40].

The finding that 70% of participants adequately hung their ITNs over the bed and tucked them under the mattress is in line with other studies that highlight the importance of correct installation and usage for effective protection. However, the 19.6% who lacked proper facilities to hang the ITNs is concerning, as previous literature has emphasized that the absence of proper infrastructure can lead to improper use, diminishing the net's effectiveness [41].

The study identified several barriers to consistent ITN use, including discomfort, heat, and structural challenges such as the inability to properly hang the nets. These are well-documented in the literature. According to Manu et al., [17] found that in many settings, the physical discomfort of using ITNs, especially during hot weather, was one of the most common reasons for non-use. The report by Doe et al., [34] also highlighted the challenges of ITN usage in hot climates, where individuals are reluctant to use the nets due to perceived discomfort. This finding aligns with the 76.7% of participants in this study who cited discomfort due to heat.

Additionally, the study found that 29.4% of participants reported the absence of proper facilities to hang ITNs, and 34.2% found the daily routine of raising the net each morning discouraging. These logistical barriers are common in the literature and have been shown to hinder regular use [17]. The findings on structural issues—such as inadequate space to hang the ITN or the routine of moving the net—are particularly important, as they highlight the need for more than just the provision of ITNs. They also emphasize the necessity of improving living conditions and providing education on how to use ITNs effectively.

The study showed that education level, employment status, and household income were significant determinants of the insecticide-treated mosquito net (ITN) usage among pregnant women. The study showed that education consistently emerges as a determinant of ITN utilization in several studies. Higher education levels are often associated with increased health literacy, awareness of malaria prevention strategies, and better adoption of ITNs. For instance, studies by Hill et al. [42] in Sub-Saharan Africa emphasize that educated women

are more likely to understand the importance of ITNs and adhere to their use. The findings that basic, secondary, and tertiary education levels are associated with reduced odds of non-usage compared to no formal education align with these patterns [43,44]. However, the observation of a less pronounced protective effect for tertiary education compared to lower levels diverges slightly from global trends. This could indicate that foundational education levels are sufficient to instill malaria prevention knowledge, with higher education providing diminishing returns in awareness specific to ITNs.

Employment status also plays a complex role. The reduced odds of non-usage among self-employed and unemployed women compared to formally employed individuals diverge from the conventional assumption that employment facilitates health resource access. This divergence may reflect the time flexibility of self-employed women, who may prioritize household health measures, including ITN use, over work commitments. Similarly, unemployed women may have more opportunities to engage with community health campaigns or receive ITNs through targeted programs. These findings echo patterns reported in regions where formal employment is associated with time constraints, reducing the likelihood of engaging in preventive health behaviors.

Household income shows a consistent relationship with ITN usage, as higher-income households generally have better access to health resources. Higher income reduces financial barriers to purchasing ITNs when they are not freely distributed and supports better living conditions, which may reduce exposure to malaria. However, the significant reduction in non-usage odds observed across all higher-income brackets suggests that even modest increases in income can make a substantial difference in ITN accessibility and utilization. This pattern is consistent with findings from Polec et al., [45], who observed that small increases in household income improved malaria prevention practices in resource-limited settings.

This study's strengths include its high participation rate, robust sample size, and thorough quantitative analysis of factors influencing ITN utilization among pregnant women in Tamale Metropolis. The focus on socio-demographic determinants provides valuable insights into how education, employment, and income levels affect ITN

usage. The study also benefits from clear data collection methods and statistical analysis that allow for meaningful comparisons between different socio-economic groups. However, the study has limitations, such as its reliance on self-reported data, which may introduce biases regarding ITN use and awareness. The cross-sectional design limits the ability to establish causal relationships, and the findings may not be generalizable to other regions outside Tamale Metropolis due to cultural and geographical differences. Additionally, the study does not assess the long-term impact of ITN usage on malaria prevention outcomes, restricting its scope in evaluating the broader effectiveness of malaria control programs.

5. CONCLUSION

This study highlights the significant role of education, employment status, and household income in determining the utilization of insecticide-treated mosquito nets (ITNs) among pregnant women in the Tamale Metropolis. Despite high awareness and ownership of ITNs, consistent usage remains low due to discomfort from heat, lack of proper hanging facilities, and logistical challenges. These barriers, alongside socio-economic disparities, emphasize the need for targeted interventions to improve the accessibility and usability of ITNs. Efforts should focus on enhancing educational campaigns, addressing environmental discomfort, and ensuring the provision of adequate infrastructure for ITN installation. The study underscores the importance of a multifaceted approach, combining awareness, resources, and community engagement, to increase the effectiveness of malaria prevention strategies in the region.

6. RECOMMENDATIONS

Pregnant women should be encouraged to use ITNs consistently by overcoming barriers like discomfort. Education on proper ITN usage and managing heat discomfort can improve adherence. Women should be urged to access free ITNs through ANC services and other distribution programs to ensure continuous protection against malaria.

The Ministry of Health should ensure better access to ITNs, particularly in rural areas, and focus on overcoming infrastructural challenges. Enhanced educational campaigns should target pregnant women, emphasizing the importance of

ITNs and their proper use. Special attention should be given to addressing discomfort issues, such as designing more breathable nets or providing guidelines to manage heat during sleep.

Midwives should be trained to educate pregnant women on the importance of ITN use, proper installation, and maintenance. They should actively advocate for using ITNs consistently during ANC visits and help overcome heat discomfort and installation challenges.

Future research should explore socio-cultural factors that influence ITN use and identify strategies to overcome barriers related to discomfort and infrastructure. Longitudinal studies assessing the impact of ITN usage on maternal and child health outcomes would provide valuable data, while investigating alternative malaria prevention methods for pregnant women could offer additional insights.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology.

Details of the AI usage are given below:

1. Authors at this moment declare that generative AI (ChatGPT) has been used during manuscript editing(grammar).

DATA AVAILABILITY

Data used to support this study are available from the corresponding author upon request.

FUNDING STATEMENT

The authors did not receive specific funding for this work.

ETHICAL APPROVAL AND CONSENT

The study was conducted in adherence to the ethical principles outlined in the Declaration of Helsinki. These included informed consent, confidentiality, non-discrimination, inclusivity, risk minimization, cultural sensitivity, and ethical

training. Participants provided written informed consent, and personal data was kept secret. The recruiting procedure was fair, and the interviews were brief, polite, and unobtrusive. Cultural sensitivity was maintained, and data collectors received training in ethical practices. The study upheld the participants' rights, dignity, and safety.

ACKNOWLEDGEMENT

We thank all the study participants who availed themselves of being part of the study at will.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- World Health Organization. World malaria report 2023. World Health Organization; 2023.
- Wafula ST, Habermann T, Franke MA, et al. What are the pathways between poverty and malaria in sub-Saharan Africa? A systematic review of mediation studies. *Infect Dis Poverty*. 2023;12:13–30.
- World Health Organization. Insecticide-treated nets to reduce the risk of malaria in pregnant women. World Health Organ.
- Mueller I, Tronick E. Early life exposure to violence: Developmental consequences on brain and behavior. *Front Behav Neurosci*. 2019;13:156.
- Mueller DH, Wiseman V, Bakusa D, et al. Cost-effectiveness analysis of insecticide-treated net distribution as part of the Togo Integrated Child Health Campaign. *Malar J*. 2008;7:1–7.
- World Health Organization. The Roll Back Malaria strategy for improving access to treatment through home management of malaria. World Health Organization; 2005.
- World Bank. Rolling Back Malaria: The World Bank Global Strategy & Booster Program. World Bank Publications; 2005.
- Asumah MN, Akugri FA, Akanlu P, et al. Utilization of insecticides treated mosquito bed nets among pregnant women in Kassena-Nankana East municipality in the upper east region of Ghana. *Public Heal Toxicol*. 2021;1:1–11.
- Alhassan Y, Dwomoh D, Amuasi SA, et al. malaria prevalence among women of reproductive age in Ghana: implication for malaria control and elimination. *Malar J*. 2022;21:120.
- Bawuah A, Ampaw S. Ownership and use of insecticide-treated nets under Ghana's National Malaria Control Program: What are the correlates? *Trop Med Int Heal*. 2021;26:1593–1608.
- Lukwa AT, Atta-Obeng C, Kaseke TW, et al. Malaria finance, household mosquito net ownership & use, intermittent preventive treatment in pregnancy (IPTp) in Ghana; a socioeconomic comparative analysis. *Health (Irvine Calif)*. 2020;20:21.
- Budu E, Okyere J, Mensah F, et al. Inequalities in the use of insecticide-treated nets by pregnant women in Ghana, 2011 and 2017. *Malar J*. 2022;21:376.
- Forty J, Keetile M. Patterns and correlates of ownership and utilization of insecticide-treated bed-nets for malaria control among women of reproductive age (15–49 years) in Malawi. *J Biosoc Sci*. 2022;54:269–278.
- Pons-Duran C, Llach M, Sacoar C, et al. Coverage of intermittent preventive treatment of malaria in pregnancy in four sub-Saharan countries: findings from household surveys. *Int J Epidemiol*. 2021;50:550–559.
- World Health Organization. Updated WHO recommendations for malaria chemoprevention among children and pregnant women.
- Kabuya J-BB, Ippolito MM, Sikalima J, et al. Safety and efficacy of intermittent presumptive treatment with sulfadoxine-pyrimethamine using rapid diagnostic test screening and treatment with dihydroartemisinin-piperaquine at the first antenatal care visit (IPTp-SP+): study protocol for a randomized. *Trials*. 2021; 22:1–12.
- Manu G, Boamah-Kaali EA, Febir LG, et al. Low utilization of insecticide-treated bed net among pregnant women in the middle belt of Ghana. *Malar Res Treat*; 2017.
- Núñez L, Skjefte M, Asamoah OE, et al. Successful implementation of ITN distribution through health facilities in Ghana. *Malar J*. 2023;22:224.
- Koenker H, Kumoji EK, Erskine M, et al. Reported reasons for non-use of insecticide-treated nets in large national household surveys, 2009–2021. *Malar J*. 2023;22:61.

20. Kviz FJ. Conducting health research: Principles, process, and methods. Sage Publications; 2019.
21. Appiah F, Salihu T, Fenteng JOD, et al. Postnatal care utilisation among women in rural Ghana: analysis of 2014 Ghana demographic and health survey. *BMC Pregnancy Childbirth*. 2021;21:1–9.
22. Nyavor KD, Kweku M, Agbemafle I, et al. Assessing the ownership, usage and knowledge of insecticide treated nets (ITNs) in malaria prevention in the hohoe municipality, Ghana. *Pan Afr Med J*; 28.
23. Paul P, Mondal D. Maternal experience of intimate partner violence and its association with morbidity and mortality of children: Evidence from India. *PLoS One*. 2020;15:e0232454.
24. Kanmiki EW, Awoonor-Williams JK, Phillips JF, et al. Socio-economic and demographic disparities in ownership and use of insecticide-treated bed nets for preventing malaria among rural reproductive-aged women in northern Ghana. *PLoS One*. 2019;14:e0211365.
25. Saha A, Sarker M, Kabir M, et al. Knowledge, attitudes, and practices regarding malaria control among the slash and burn cultivators in Rangamati Hill tracts of Bangladesh. *Malar J*. 2019;18:1–9.
26. Osman ASA, Ahmed MA, Mohamed MH, et al. Factors Affecting the Utilization of Antenatal Care among Married Women of Reproductive Age in Merca, Lower Shebelle, Somalia. *J Biosci Med*. 2022;10:165–175.
27. Pryce J, Richardson M, Lengeler C. Insecticide-treated nets for preventing malaria. *Cochrane Database Syst Rev*.
28. Lindblade KA, Mwandama D, Mzilahowa T, et al. A cohort study of the effectiveness of insecticide-treated bed nets to prevent malaria in an area of moderate pyrethroid resistance, Malawi. *Malar J*. 2015;14:1–15.
29. Wangdi K, Furuya-Kanamori L, Clark J, et al. Comparative effectiveness of malaria prevention measures: a systematic review and network meta-analysis. *Parasit Vectors*. 2018;11:1–13.
30. Ahorlu CS, Adongo P, Koenker H, et al. Understanding the gap between access and use: A qualitative study on barriers and facilitators to insecticide-treated net use in Ghana. *Malar J*. 2019;18:1–13.
31. Scott J, Kanyangarara M, Nhama A, et al. Factors associated with use of insecticide-treated net for malaria prevention in Manica District, Mozambique: a community-based cross-sectional survey. *Malar J*. 2021;20:200.
32. Nishimwe C, Mchunu GG, Mukamsoni D. Community-based maternal and newborn interventions in Africa: systematic review. *J Clin Nurs*. 2021;30:2514–2539.
33. Benito E, Suthar P, Banda G, et al. Perception of caregivers on ITNs utilisation and its effectiveness among children under 5 years of age in Chikwawa district, Malawi. *Malar J*. 2024;23:341.
34. Doe PF, Druye AA, Azu TD, et al. Ownership and usage of insecticide-treated nets in Ghana: A scoping review of facilitators and barriers. *Malar J*. 2024;23:238.
35. Tizifa TA, Kabaghe AN, McCann RS, et al. Prevention efforts for malaria. *Curr Trop Med Reports*. 2018;5:41–50.
36. Sserwanja Q, Mutisya LM, Musaba MW. Exposure to different types of mass media and timing of antenatal care initiation: Insights from the 2016 Uganda Demographic and Health Survey. *BMC Womens Health*. 2022;22:1–8.
37. Khanal V, Adhikari M, Karkee R, et al. Factors associated with the utilisation of postnatal care services among the mothers of Nepal: analysis of Nepal demographic and health survey 2011. *BMC Womens Health*. 2014;14:1–13.
38. Kanmiki EW, Awoonor-Williams JK, Phillips JF, et al. Predictors of mosquito net use in Ghana. *New York*. 2014;14:1–6.
39. Munisi DZ, Nyundo AA, Mpondo BC. Knowledge, attitude and practice towards malaria among symptomatic patients attending Tumbi Referral Hospital: A cross-sectional study. *PLoS One*. 2019;14:e0220501.
40. DePina AJ, Dia AK, Martins A de AS, et al. Knowledge, attitudes and practices about malaria in Cabo Verde: a country in the pre-elimination context. *BMC Public Health*. 2019;19:1–14.
41. Lengeler C. Insecticide-treated bed nets and curtains for preventing malaria. *Cochrane Database of Syst Rev*; 2008. Issue 2 Art No CD000363.
42. Hill J, Hoyt J, van Eijk AM, et al. Factors affecting the delivery, access, and use of interventions to prevent malaria in pregnancy in sub-Saharan Africa: a systematic review and meta-analysis. *PLoS Med*. 2013;10:e1001488.

43. Eshetu B, Bekele H, Debella A, et al. Insecticide-treated net utilization and associated factors among pregnant women in Ethiopia: A systematic review and meta-analysis. *Front Glob Women's Heal.* 2023;4:1147583.
44. Onyinyechi OM, Mohd Nazan AIN, Ismail S. Effectiveness of health education interventions to improve malaria knowledge and insecticide-treated nets usage among populations of sub-Saharan Africa: systematic review and meta-analysis. *Front Public Heal.* 2023;11:1217052.
45. Polec LA, Petkovic J, Welch V, et al. Strategies to increase the ownership and use of insecticide-treated bednets to prevent malaria. *Campbell Syst Rev.* 2015;11:1–127.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/127714>