

Asian Journal of Cardiology Research

Volume 8, Issue 1, Page 35-57, 2025; Article no.AJCR.129229

Impact of Aerobic Exercise on Obesity among Female Undergraduates of Ladoke Akintola University of Technology, Ogbomoso, Nigeria

A. M. Adesola ^{a*}, F. O. Akinwumi ^a, O. D. Ogunleye ^a and S. O. Hammed ^a

^a Physiology Department, Faculty of Basic Medical Sciences, College of health Science, Ladoke Akintola University of Technology (LAUTECH) Ogbornoso, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: https://doi.org/10.9734/ajcr/2025/v8i1242

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/129229

Original Research Article

Received: 26/10/2024 Accepted: 02/01/2025 Published: 11/01/2025

ABSTRACT

Introduction: According to the World health organization, obesity is an abnormal or excessive fat accumulation which impairs health, and is caused by imbalance between calories intake and calories expended. In the 21st century, overweight and obesity were only problems of developed countries which have increased due to sedentary life styles, behavioral changes, and genetic modifications. Obesity is dominating as a global epidemic as it increases risk of human life which is directly related to some diseases such as hypertension, cardiovascular diseases, musculoskeletal disorders, certain cancer types and diabetes. Hence the need for effective solution for obesity has led to the recommendation of aerobic exercises that help in weight reduction via increasing the body's ability to utilize oxygen efficiently. This study aimed to evaluate the effect of aerobic exercise

Cite as: Adesola, A. M., F. O. Akinwumi, O. D. Ogunleye, and S. O. Hammed. 2025. "Impact of Aerobic Exercise on Obesity Among Female Undergraduates of Ladoke Akintola University of Technology, Ogbomoso, Nigeria". Asian Journal of Cardiology Research 8 (1):35-57. https://doi.org/10.9734/ajcr/2025/v8i1242.

^{*}Corresponding author: Email: amadesola@lautech.edu.ng;

on obesity and weight management body in female obese subjects (undergraduates) of Ladoke Akintola University of Technology Ogbomoso.

Forty (40) female obese subjects were randomly selected for this study. Subjects were undergraduate students from Ladoke Akintola University of Technology (LAUTECH), within the age range of about (18-30) years. All subjects signed the written consent form prior to the study and health status questionnaire to gather demographic information and data on lifestyle for the study. The subjects were randomly grouped into 2 (n=20); Control group that did not perform any exercise and experimental group that performed aerobic exercise with 60% exercise intensity according to the 'Karovonen' formulae for 50minutes daily which lasted for 3 days and for 50minutes weekly which lasted for 8weeks.

Pre- and post- anthropometric measurement (body weight, height, body mass index, circumferential and skinfold measurements), cardiovascular characteristics (heart rate and blood pressure), and biochemical assays [blood glucose level, triglyceride, cholesterol, low density lipo-protein and high density lipo-protein] were done on both control and experimental subjects to assess the effect of aerobic exercise on obesity. Statistical analysis of data was done using SPSS to determine the mean and the standard deviation, T-test was done to determine P-value, which was set as P<0.05 which determine the level of significant in the two groups.

The control group maintains its average body weight of 80.80 kg, height of 1.622m and BMI of 30.75; blood pressure and heart rate of control subjects of 134.1/73.20mmHg and 72beat/minute; average blood glucose level of (112.6 \pm 12.04); lipid profile levels, circumferential and skinfold measurement throughout the study. However, experimental group showed significant reduction in: the average body weight and BMI of experimental subjects from 86.25kg and 33.00kg/m² to 77.65and 29.45 respectively (p= 0.001 and 0.014), Blood pressure from 133.1/73.40mmHg to 115.1/64.25mmHg (p= 0.017), the average blood glucose level from 111.0/dl to 80.70g/dl (p= 0.001), [LDL, TRIG, and Cholesterol levels] (p= 0.001), circumferential and skinfold measurement and of certain body parts at the end of the aerobic exercise and when compared to control subjects (p<0.05).

Conclusion: Aerobic exercise significantly reduces body excess compositions, blood glucose level, body mass index, blood pressure lipid profile except for high density lipo-protein, showing that aerobic exercise is of great health benefit.

Keywords: Aerobic exercise; heart rate and blood pressure; body compositions; obesity.

1. INTRODUCTION

Obesity is a condition characterized bv accumulation of excessive amount of body fat beyond the biological need of an individual. Obesity is a principle public health concern ranked as the fifth foremost reasons for global mortality (WHO, 2016). From the last 40years, obesity is dominating as a global epidemic and is raising non communicable diseases such as type 2 diabetes, hypertension, metabolic syndrome, certain type of cancer and cardiovascular diseases that threaten human health globally (Lee et al.; 2013). In 2017 more than 15% of deaths were related to obesity in many developing countries such as Eastern Europe, Central Asia, North Africa and western Africa and Latin America. Obesity is a manifestation of a positive energy balance that has been sustained over an extended period of time. Although, human genome has not changed substantially: but the rise in obesity most likely reflects changes in the environment and/or behavior. In

the 20st century overweight and obesity were only problems of developed countries but in the 21st century, these are increased in lower developed and developing countries due to the popularity of junk foods, changing lifestyles, increasing incomes, urbanization and aggressive marketing (Ng and Popkin, 2012). Mechanization has impinged upon our modes of living in diverse ways and energy expenditure required for daily living has consequently declined. A recent study showed that in the US, daily energy expenditure due to work related physical exercise has decreased by more than 100 kcal during last 50 years in both men and women, and this reduction is associated with the increase in mean body weight during this time frame (Church et al., 2005). Similar trends have also been observed in Finland, where daily energy expenditure during work reportedly decreased by more than 50 kcal between 1982 and 1992 (Fogelholm et al., 1996). Substantial reductions in daily energy expenditure have also occurred in developing countries such as China and Brazil, which have the highest absolute and relative rates of decline in total energy expenditure due to restrictions of movement at work (Ng and Popkin, 2012). Based on these reports, it is believed that the obesity epidemic has also penetrated the low-income countries, particularly in the urban areas, and will continue to spread for the foreseeable future (Prentice, 2006).

Thus, there is a need to reverse this modern epidemic and public health actions to reduce obesity have mostly advised obese individuals to eat healthier and to exercise more. However, these approaches are failing as not a single country has succeeded in reducing obesity rates in the past 30 years (Ng et al., 2014). It is yet to be determined whether this is due to failure to restrict energy intake or to maintain high levels of energy expenditure (Blair et al., 2013; Hill and Peters, 2013; Luke and Cooper, 2013). Domestic mechanization of daily tasks (use of labor-saving devices such as washing machines and dishwashers) has reduced energy expenditure over the years (Archer et al., 2013). Domestic mechanization has also contributed to increased sedentariness, as time spent in house work has been replaced by sedentary activities such as watching television, and the use of smart phones and computers (Archer et al., 2013). Passive transportation has also been linked to be a major cause of excessive weight gain (Thorp et al., 2011; Rhodes et al., 2012; McCormack et al., 2014). Participation in leisure-time physical activity (LTPA) on a secular basis with unhealthy eating habit are not enough to offset sedentary behavior (Ng and Popkin, 2012). Hence the need for effective solution for obesity has led to the recommendation of aerobic exercises that help in weight reduction. Aerobic exercise, a type of physical activity involves the continuous and rhythmic movement of large muscle groups to improve the efficiency of the cardiovascular and respiratory systems in delivering oxygen to the body (Stasiulis et al., 2004). Types of aerobic exercise include; walking, jogging, swimming, skipping and cycling. These physical activities enable rapid breathing, rapid heartbeat and sweating to increase the body's ability to utilize oxygen efficiently. In this study, the role of aerobic exercise in obesity and weight management was explored.

2. MATERIALS AND METHODS

2.1 Materials and Chemicals

Yoga mat, bicycle ergometer, skipping rope, treadmill, multi-gym equipment, hand grip

exerciser, Avery-weighing machine, measuring tape, meter rule, automatic blood pressure monitor, centrifuge machine, micro pipette, test tubes, needle and syringe, hand gloves, methylated spirit and cotton wool.

2.2 Study Area

This research study was carried out in the sport and exercise gym of the department of physiology, faculty of Basic Medical Sciences, Ladoke Akintola University of Technology.

2.3 Subjects

Forty (40) female obese subjects were randomly selected for this studv. Subiects were undergraduate students from Ladoke Akintola University of Technology (LAUTECH), within the age range of about (18-30) years. All subjects were aware of the risk and requirement of participating in the study and they all signed the written consent form prior to the study. The subjects filled health status questionnaire to gather demographic information, as well as data on lifestyle for the study. With the use of health status questionnaire and physical examination, individuals with underlying heart, lungs, and systemic diseases were excluded. Alcoholics and cigarette smokers were also excluded.

2.4 Subject Grouping

Forty obese female subjects were randomly grouped into 2 (n=20):

Group 1: Control group: The subjects in this group did not perform any exercise.

Group 2: Experimental group: The subjects in this group performed aerobic exercise with 60% exercise intensity according to the 'Karovonen' formulae for 50minutes daily which lasted for 3 days (Willis et al., 2012) and for 50minutes weekly which lasted for 8weeks (Aghaei et al., 2018). The subjects were dressed in sports kits so as to aid easiness of performing the exercise.

Upon their arrival to the laboratory, they were made to sit for 10minutes to normalize their heart rate and blood pressure, after which the physical characteristics and the cardiovascular characteristics of the experimental and control subjects were taken before the experimental subject began exercising.

2.5 Aerobic Exercise Types

The aerobic exercises carried out in this study include: Yoga, cycling, skipping, running and grip strength training. The subjects were monitored when carrying out the exercise to ensure adherence to the exercise regimen.

2.5.1 Pre- and post- anthropometric measurement of subjects

1. Body Mass Index (BMI)

Body Mass Index (BMI), is a measure of body fat based on height and weight is calculated by the use of this formula (WHO, 2008):

BMI=weight (kg)/ height² (m)

(i) Height Measurement

The standing position of the subjects was measured in meter on a Stadiometer [accuracy of 0.01m] with bare feet, person looking straight ahead, positioning the head in the Frankfort plane, with relaxed shoulders, the scapula, buttocks and heels should be touching the wall, arms on each side of the body, legs straight and together to face.

(ii) Body Weight Measurement

Body weight of the subjects (in kg) was measured using weighing scale with a precision of ± 0.1 kg. The subject was allowed to stand on the weighing scale with bare feet and face up for full body weight measurement.

2. Circumferential Measurement

The burst, chest, abdomen, hip, thigh (right and left), calf (right and left), upper arm (right and left), lower arm (right and left) was measured in inches with a non-elastic tape rule.

3. Skinfold Measurement

The thickness of skinfold of the biceps (right and left), triceps (right and left), chest, abdomen, bi- iliac, sub-scapular, inner thigh (left and right) and calf (left and right) was measured in millimeter with a digital caliper.

Anthropometric measurement like circumferential measurement and skinfold measurement were conducted for both experimental and control for pre-test and post-test.

2.5.2 Calculation of exercise intensity in participants

Exercise intensity and target heart rate was calculated using the 'Karovonen' formula (Karovonen et al., 1957):

Target Heart Rate = [(max HR - resting HR) × %Intensity] + resting HR. (Exercise intensity= 60%)

2.5.3 Cardiovascular characteristics

Pre- and post- Heart rate and blood pressure was measured using a digital sphygmomanometer (Omron Automatic Digital Sphyg.) while the subject is in sitting position and cuff tied round the arm at 4cm above the cubital fossa.

2.6 Sample Collection

A 2ml of blood were collected from both control and experimental subjects before performing exercise for biochemical determination (pre-test). blood samples were collected The into heparinized sample bottles, centrifuged at 3000R for 10minutes to obtain plasma that was collected into plain tubes for biochemical glucose [blood] assavs level, triglyceride, cholesterol, low density lipo-protein and high density lipo-protein]. After the last day of the exercise, biochemical analysis was done on the subject blood samples (post-test) similar to that of pre-test.

2.7 Method of Statistical Analysis

Data from all participants were analyzed using SPSS and were recorded as Means \pm SD; while the T-test for paired samples was applied to test for differences on data obtained from aerobics exercise. The statistical level of significance was accepted at p < 0.05.

3. RESULTS

1. Pre-test and post-test of physical characteristics (Table 1a-d): The control subjects maintain the average body weight of 80.80 kg, height of 1.622m and BMI of 30.75 throughout the study. However, table significant shows reduction 2 а average body weight and in the BMI of experimental subjects from 86.25kg and 33.00kg/m² to 77.65and 29.45 respectively (P= 0.001 and 0.014) at the end of the aerobic exercise. Although, there was no significant difference in the BMI of control and experimental subjects, but there was a significant reduction in the average body weight of experimental subjects compared to control subjects (p=0.0364)

- 2. Pre-test and post-test of cardiovascular function (Table 2a-d): Blood pressure and heart rate of control subjects remain constant at 134.1/73.20mmHg and 72beat/minute throughout study the duration. Whereas, experimental subjects showed decreased Blood pressure from 133.1/73.40mmHg to 115.1/64.25mmHg (P<0.05) (P= 0.017) at the end of the aerobic exercise and constant heart rate of 72.40beat/minute. Compared to control subjects, experimental subjects showed decreased Blood pressure (P= 0.0032) at the end of the aerobic exercise.
- **3. Pre-** and post-glucose test (Table 3a-d): There was no significant difference in the average blood glucose level of control subjects before (112.6±12.04) and after (110.2± 12.57) the study. On the contrary, the average blood glucose level of experimental subjects significantly decreased from 111.0/dl to 80.70g/dl at the end of the aerobic exercise and when compared with control subjects, p= 0.001.
- 4. Pre- and post- training lipid profile (Table 4a-d): The control subjects show no significant difference in the lipid profile before and after the study. However, experimental subjects show significant increase in HDL at the end of the aerobic exercise (0.015) and when compared to control subjects (0.001), as well as a significant decrease in LDL, TRIG, and Cholesterol levels at the end of the aerobic exercise (p= 0.001) and when compared to control subjects (p= 0.001).
- post-5. Preand circumferential measurement (Table 5a-d): The control subjects show no significant difference in circumferential measurement the throughout the study duration. Nonetheless, experimental subjects show significant reduction in circumferential measurement of certain body parts at the end of the aerobic exercise and compared to control subjects (p<0.05).

6. Pre- and post- skinfold measurement (Table 6a-d): The control subjects show no significant difference in the skinfold measurement throughout the study duration. However, experimental subjects show significant reduction in skinfold measurement of body parts with high subcutaneous tissue at the end of the aerobic exercise and compared to control subjects (p<0.05).

4. DISCUSSION

This study showed that aerobic exercise significantly decreased body mass index (BMI) and body composition measurement which include circumferential measurement, skinfold measurement, weight, height, heart rate and blood pressure in the experimental subjects compare to the control subjects that did not partake in the intervention program (aerobic exercise). The decrease could have occurred due to reduction in body fat mass and decreased body weight and body mass index (BMI) caused by exercise this is in line with study carried out by (Stasiulis et al, 2010). Previous study has also shown that aerobic exercise reduces body mass index, body fat mass, and body mass in which aerobic exercise causes the capability of the body to use fat as a substrate to increase the total fat oxidation (Chiu et al, 2017). Aerobic exercise lower blood pressure by improving circulation, strengthening the heart, and reducing stress exercise improves circulation which can lower blood pressure and heart rate. Exercise triggers nitric oxide production to promote cardiovascular health via inducing vasodilation and reducing atherosclerosis (Li et al., 2021). Aerobic exercise activates lipoprotein lipase and increased lipoprotein lipase activities play an important role in reducing insulin resistance during aerobic exercise in line with the study carried out (Fenkcis et by al.,2006). In obese individual aerobic exercise plays an s important role in reducing body fat by enhancing oxygen utilization by combining with glucose to release energy in form of adenosine triphosphate when glucose is being exhausted, the body make use of fat through gluconeogenesis process to generate energy which helps in reducing body fat in line with the study carried out by (Plowman and Smith. 2007)

| S/N | Age | Weight | Height | BMI (Ka/M ²) | S/N | Age | Weight | Height | BMI |
|-------|--------------|---------------|--------|-----------------------------|-------|--------|--------|--------|-----------|
| 4 | <u>(yrs)</u> | (K <u>G</u>) | (III) | (r.g/wi-) | 4 | (yrs) | (Kg) | (11) | (Kg/IVI-) |
| 1 | 18 | 85 | 1.70 | 29.41 | 1 | 18 | 85 | 1.70 | 29.41 |
| 2 | 24 | 75 | 1.57 | 30.42 | 2 | 24 | 75 | 1.57 | 30.42 |
| 3 | 24 | 91 | 1.61 | 35.10 | 3 | 24 | 91 | 1.61 | 35.10 |
| 4 | 18 | 82 | 1.55 | 34.13 | 4 | 18 | 82 | 1.55 | 34.13 |
| 5 | 19 | 72 | 1.62 | 27.43 | 5 | 19 | 72 | 1.62 | 27.43 |
| 6 | 20 | 80 | 1.65 | 29.38 | 6 | 20 | 80 | 1.65 | 29.38 |
| 7 | 27 | 75 | 1.60 | 29.29 | 7 | 27 | 75 | 1.60 | 29.29 |
| 8 | 19 | 73 | 1.54 | 30.74 | 8 | 19 | 73 | 1.54 | 30.74 |
| 9 | 25 | 78 | 1.67 | 27.96 | 9 | 25 | 78 | 1.67 | 27.96 |
| 10 | 20 | 85 | 1.66 | 30.84 | 10 | 20 | 85 | 1.66 | 30.84 |
| 11 | 20 | 84 | 1.56 | 34.51 | 11 | 20 | 84 | 1.56 | 34.51 |
| 12 | 25 | 79 | 1.67 | 28.32 | 12 | 25 | 79 | 1.67 | 28.32 |
| 13 | 19 | 85 | 1.54 | 35.84 | 13 | 19 | 85 | 1.54 | 35.84 |
| 14 | 27 | 80 | 1.69 | 28.01 | 14 | 27 | 80 | 1.69 | 28.01 |
| 15 | 20 | 80 | 1.65 | 28.68 | 15 | 20 | 80 | 1.65 | 28.68 |
| 16 | 19 | 87 | 1.62 | 33.15 | 16 | 19 | 87 | 1.62 | 33.15 |
| 17 | 18 | 82 | 1.65 | 30.11 | 17 | 18 | 82 | 1.65 | 30.11 |
| 18 | 24 | 90 | 1.61 | 34.63 | 18 | 24 | 90 | 1.61 | 34.63 |
| 19 | 24 | 73 | 1.57 | 29.61 | 19 | 24 | 73 | 1.57 | 29.61 |
| 20 | 18 | 80 | 1.70 | 27.68 | 20 | 18 | 80 | 1.70 | 27.68 |
| Mean± | 21.20± | 80.80± | 1.622± | 30.75± | Mean± | 21.20± | 80.80± | 1.622± | 30.75± |
| SD | 3.139 | 5.49 | 0.0532 | 2.776 | SD | 3.139 | 5.49 | 0.0532 | 2.776 |

Table 1a. Pre-test of physical characteristics of control subjects

Table 1b. Post-test of physical characteristics of control subjects

| S/N | Age | Weight | Height | BMI | S/N | Age | Weight | Height | |
|-------|--------|-------------|------------|--------|-------|--------|--------------|--------|--------|
| 4 | | <u>(Kg)</u> | <u>(m)</u> | | | | <u>(K</u> g) | (III) | |
| 1 | 27 | 88 | 1.67 | 31.56 | 1 | 27 | 80 | 1.67 | 28.68 |
| 2 | 22 | 90 | 1.51 | 39.47 | 2 | 22 | 82 | 1.51 | 35.96 |
| 3 | 24 | 85 | 1.58 | 34.04 | 3 | 24 | 70 | 1.58 | 28.04 |
| 4 | 19 | 86 | 1.67 | 30.83 | 4 | 19 | 80 | 1.67 | 28.68 |
| 5 | 19 | 90 | 1.66 | 32.66 | 5 | 19 | 80 | 1.66 | 30.48 |
| 6 | 24 | 81 | 1.61 | 31.24 | 6 | 24 | 75 | 1.61 | 28.93 |
| 7 | 27 | 90 | 1.66 | 32.66 | 7 | 27 | 80 | 1.66 | 29.03 |
| 8 | 18 | 75 | 1.56 | 30.81 | 8 | 18 | 70 | 1.56 | 28.76 |
| 9 | 19 | 80 | 1.59 | 31.64 | 9 | 19 | 75 | 1.59 | 29.66 |
| 10 | 24 | 85 | 1.67 | 30.47 | 10 | 24 | 75 | 1.67 | 26.89 |
| 11 | 28 | 82 | 1.62 | 31.24 | 11 | 28 | 76 | 1.62 | 28.95 |
| 12 | 23 | 79 | 1.56 | 32.46 | 12 | 23 | 70 | 1.56 | 28.76 |
| 13 | 20 | 85 | 1.58 | 34.04 | 13 | 20 | 75 | 1.58 | 30.04 |
| 14 | 27 | 90 | 1.66 | 32.66 | 14 | 27 | 82 | 1.66 | 29.75 |
| 15 | 24 | 81 | 1.61 | 31.24 | 15 | 24 | 70 | 1.61 | 27.00 |
| 16 | 22 | 90 | 1.62 | 34.29 | 16 | 22 | 85 | 1.62 | 32.38 |
| 17 | 26 | 95 | 1.67 | 34.06 | 17 | 26 | 87 | 1.67 | 31.19 |
| 18 | 29 | 98 | 1.68 | 34.72 | 18 | 29 | 87 | 1.68 | 30.82 |
| 19 | 29 | 90 | 1.51 | 39.41 | 19 | 29 | 80 | 1.51 | 28.56 |
| 20 | 27 | 85 | 1.67 | 30.47 | 20 | 27 | 74 | 1.67 | 26.53 |
| Mean± | 23.90± | 86.25± | 1.618± | 33.00± | Mean± | 23.90± | 77.65± | 1.618± | 29.45± |
| SD | 3.567 | 5.618 | 0.0545 | 2.586 | SD | 3.567 | 5.489*# | 0.0545 | 2.104* |

Table 1c. Pre- test physical characteristic of experimental subjects

Table 1d. Post- test physical characteristic of experimental subjects

Data were presented as mean ± standard deviation (SD); *= Statistical significance at p<0.05 when compared with pre-test of experimental subjects; #= Statistical significance at p<0.05 when compared with post-test of control subjects

| S/N | Systolic Blood Pressure (mmHg) | Diastolic Blood Pressure (mmHg) | Resting heart Rate (beat/minutes) | S/N | Systolic Blood Pressure (mmHg) | Diastolic Blood Pressure (mmHg) | Resting heart Rate (beat/minutes) |
|------|--------------------------------------|---------------------------------------|---|--------|--------------------------------------|---------------------------------------|---|
| 1 | 135 | 75 | 72 | 1 | 135 | 75 | 72 |
| 2 | 132 | 72 | 72 | 2 | 132 | 72 | 72 |
| 3 | 137 | 70 | 72 | 3 | 137 | 70 | 72 |
| 4 | 138 | 75 | 72 | 4 | 138 | 75 | 72 |
| 5 | 130 | 70 | 72 | 5 | 130 | 70 | 72 |
| 6 | 136 | 74 | 72 | 6 | 136 | 74 | 72 |
| 7 | 132 | 72 | 72 | 7 | 132 | 72 | 72 |
| 8 | 138 | 78 | 72 | 8 | 138 | 78 | 72 |
| 9 | 131 | 74 | 72 | 9 | 131 | 74 | 72 |
| 10 | 130 | 72 | 72 | 10 | 130 | 72 | 72 |
| 11 | 133 | 71 | 72 | 11 | 133 | 71 | 72 |
| 12 | 137 | 74 | 72 | 12 | 137 | 74 | 72 |
| 13 | 132 | 70 | 72 | 13 | 132 | 70 | 72 |
| 14 | 136 | 78 | 72 | 14 | 136 | 78 | 72 |
| 15 | 130 | 74 | 72 | 15 | 130 | 74 | 72 |
| 16 | 130 | 72 | 72 | 16 | 130 | 72 | 72 |
| 17 | 139 | 74 | 72 | 17 | 139 | 74 | 72 |
| 18 | 138 | 79 | 72 | 18 | 138 | 79 | 72 |
| 19 | 132 | 70 | 72 | 19 | 132 | 70 | 72 |
| 20 | 136 | 70 | 72 | 20 | 136 | 70 | 72 |
| Mean | 134.1± | 73.20± | 72± | Mean±S | 134.1± | 73.20± | 72± |
| ±SD | 3.19 | 2.821 | 0.00 | D | 3.19 | 2.821 | 0.00 |

Table 2a. Pre- test characteristics of blood pressure and heartrate of female obese control subjects

Table 2b. Post- test characteristics of blood pressure and heart rate of
female obese control subjects

| S/N | Systolic Blood Pressure (mmHg) | Diastolic Blood Pressure (mmHg) | Resting heart Rate (beat/minutes) | S/N | Systolic Blood Pressure (mmHg) | Diastolic Blood Pressure (mmHg) | Resting heart Rate (beat/minutes) |
|-------|---|---------------------------------------|---|-------|---|--|--|
| 1 | 135 | 73 | 72 | 1 | 110 | 60 | 72 |
| 2 | 130 | 72 | 72 | 2 | 116 | 62 | 72 |
| 3 | 133 | 72 | 79 | 3 | 117 | 65 | 72 |
| 4 | 138 | 75 | 72 | 4 | 112 | 60 | 72 |
| 5 | 132 | 70 | 72 | 5 | 118 | 65 | 72 |
| 6 | 136 | 77 | 72 | 6 | 123 | 63 | 72 |
| 7 | 137 | 75 | 72 | 7 | 119 | 69 | 72 |
| 8 | 130 | 74 | 72 | 8 | 120 | 63 | 72 |
| 9 | 139 | 74 | 72 | 9 | 117 | 64 | 72 |
| 10 | 138 | 75 | 72 | 10 | 112 | 60 | 72 |
| 11 | 139 | 70 | 72 | 11 | 120 | 69 | 72 |
| 12 | 130 | 72 | 72 | 12 | 118 | 67 | 72 |
| 13 | 132 | 71 | 72 | 13 | 118 | 69 | 72 |
| 14 | 133 | 74 | 73 | 14 | 115 | 68 | 72 |
| 15 | 134 | 72 | 73 | 15 | 113 | 65 | 72 |
| 16 | 136 | 80 | 72 | 16 | 114 | 64 | 72 |
| 17 | 130 | 75 | 72 | 17 | 112 | 60 | 72 |
| 18 | 131 | 70 | 72 | 18 | 116 | 67 | 72 |
| 19 | 134 | 75 | 72 | 19 | 118 | 65 | 72 |
| 20 | 130 | 72 | 72 | 20 | 113 | 60 | 72 |
| Mean± | 133.1± | 73.40± | 72.40± | Mean± | 115.1± | 64.25± | 72± |
| SD | 3.233 | 2.521 | 1.56 | SD | 3.364*# | 3.226*# | 0.00 |

Table 2c. Pre- test characteristics of blood pressure and heart rate of female obese experimental subjects

Table 2d. Post- test characteristics of blood pressure and heart rate of female obese experimental subjects

Data were presented as mean \pm standard deviation (SD); *= Statistical significance at p<0.05 when compared with pre-test of experimental subjects; #= Statistical significance at p<0.05 when compared with post-test of control subjects

| S/N | Pre Glucose Level of Control Subjects (mmHg)Post Glucose Level of Control Subjects (mm | | Pre Glucose Level of Experimental Subjects | Post Glucose Level of Experimental Subjects |
|----------|--|--------------|---|--|
| | | | (mmHg) | (mmHg) |
| 1 | 115.09 | 118.66 | 118.66 | 80.53 |
| 2 | 124.25 | 114.43 | 114.43 | 75.53 |
| 3 | 118.66 | 117.40 | 117.40 | 73.19 |
| 4 | 114.43 | 109.70 | 109.70 | 83.67 |
| 5 | 117.96 | 116.82 | 116.82 | 79.20 |
| 6 | 123.09 | 124.74 | 124.74 | 79.20 |
| 7 | 114.25 | 125.63 | 125.63 | 83.67 |
| 8 | 118.66 | 113.40 | 113.40 | 73.19 |
| 9 | 84.43 | 109.70 | 109.70 | 75.19 |
| 10 | 87.96 | 116.82 | 116.82 | 90.53 |
| 11 | 93.09 | 124.74 | 124.74 | 79.20 |
| 12 | 124.25 | 125.63 | 125.63 | 83.60 |
| 13 | 122.66 | 100.40 | 100.40 | 83.90 |
| 14 | 98.43 | 109.70 | 109.70 | 75.53 |
| 15 | 110.96 | 116.82 | 116.82 | 90.53 |
| 16 | 113.09 | 84.74 | 84.74 | 85.19 |
| 17 | 124.25 | 95.63 | 95.63 | 83.67 |
| 18 | 112.66 | 100.40 | 100.40 | 75.53 |
| 19 | 115.43 | 109.70 | 109.70 | 79.20 |
| 20 | 117.96 | 96.82 | 96.82 | 85.53 |
| Mean± SD | 112.6± 12.04 | 110.2± 12.57 | 111.0±11.66 | 80.70± 5.184 ^{*#} |

Table 3. The pre and post of blood glucose level of female obese control and experimental subjects

Data were presented as mean \pm standard deviation (SD); *= Statistical significance at p<0.05 when compared with pre-test of experimental subjects; #= Statistical significance at p<0.05 when compared with post-test of control subjects

| S/N | Control | Control | Exp | Exp | Control | Control | Exp pre- | Exp |
|-------|---------|--------------|---------|----------|----------|-----------|----------|---------------------|
| · | pre-HDL | post-HUL | Pre-HDL | POST-HUL | pre-LDLP | post-LDLP | LDLP | post-LDLP |
| 1 | 65.65 | 65.65 | 59.81 | 95.27 | 119.13 | 119.13 | 129.13 | 69.35 |
| 2 | 40.79 | 40.79 | 40.79 | 92.79 | 122.28 | 122.28 | 132.28 | 65.27 |
| 3 | 36.44 | 36.44 | 38.44 | 87.85 | 124.94 | 124.94 | 126.94 | 60.15 |
| 4 | 34.11 | 34.11 | 39.11 | 78.46 | 119.09 | 119.09 | 119.09 | 75.99 |
| 5 | 38.13 | 3 8.13 | 45.13 | 78.42 | 118.23 | 118.23 | 128.23 | 67.27 |
| 6 | 37.81 | 37.81 | 37.81 | 75.27 | 131.13 | 131.13 | 131.13 | 69.35 |
| 7 | 33.79 | 33.79 | 48.79 | 76.79 | 122.28 | 122.28 | 132.28 | 66.27 |
| 8 | 36.44 | 36.44 | 40.44 | 74.85 | 124.94 | 124.94 | 124.94 | 60.15 |
| 9 | 36.07 | 36.07 | 39.07 | 74.46 | 129.09 | 129.09 | 129.09 | 68.99 |
| 10 | 37.13 | 37.13 | 34.13 | 80.42 | 121.23 | 121.23 | 131.23 | 69.27 |
| 11 | 34.81 | 34.81 | 39.81 | 97.27 | 131.13 | 131.13 | 131.13 | 69.35 |
| 12 | 34.79 | 34.79 | 42.79 | 76.79 | 122.28 | 122.28 | 127.28 | 66.27 |
| 13 | 34.44 | 34.44 | 40.44 | 79.85 | 124.94 | 124.94 | 128.94 | 60.15 |
| 14 | 32.11 | 32.11 | 32.11 | 87.46 | 139.09 | 139.09 | 139.09 | 70.99 |
| 15 | 35.13 | 35.13 | 45.13 | 78.42 | 131.23 | 131.23 | 131.23 | 79.27 |
| 16 | 36.81 | 36.81 | 39.81 | 79.27 | 131.13 | 131.13 | 131.13 | 69.35 |
| 17 | 33.79 | 33.79 | 39.79 | 86.79 | 122.28 | 122.28 | 132.28 | 76.27 |
| 18 | 33.79 | 33.79 | 40.44 | 73.85 | 124.94 | 124.94 | 127.94 | 60.15 |
| 19 | 34.11 | 34.11 | 39.11 | 94.46 | 129.09 | 129.09 | 129.09 | 80.99 |
| 20 | 35.13 | 35.13 | 54.13 | 78.42 | 131.23 | 131.23 | 131.23 | 79.27 |
| Mean± | 37.25± | 37.25± 6.966 | 37.25± | 41.85± | 126.0± | 126.0±5 | 129.7± | 69.21± |
| SD | 6.966 | | 6.96 | 6.361*# | 5.439 | .439 | 3.818 | 6.510 ^{*#} |

Table 4a. Pre- and post-training Lipid profile of obese female control and experimental subjects

Data were presented as mean ± standard deviation (SD); *= Statistical significance at p<0.05 when compared with pre-test of experimental subjects; #= Statistical significance at p<0.05 when compared with post-test of control subjects

| S/N | Control | Control | Exp | Exp | Control | Control | Exp | Exp | |
|-------|----------|-----------|----------|-----------|----------|-----------|----------|-----------|--|
| | pre-TRIG | post-TRIG | pre-TRIG | post-TRIG | pre-CHOL | post-CHOL | pre-CHOL | post-CHOL | |
| 1 | 189.27 | 189.27 | 189.27 | 124.57 | 214.48 | 214.48 | 217.48 | 190.53 | |
| 2 | 228.60 | 228.60 | 228.60 | 106.96 | 220.03 | 220.03 | 229.03 | 180.45 | |
| 3 | 260.84 | 260.84 | 261.84 | 122.44 | 241.71 | 241.71 | 237.71 | 182.49 | |
| 4 | 242.54 | 242.54 | 244.54 | 135.33 | 281.75 | 281.75 | 281.75 | 178.52 | |
| 5 | 240.01 | 240.01 | 239.01 | 145.06 | 215.38 | 215.38 | 225.38 | 170.70 | |
| 6 | 329.27 | 329.27 | 329.27 | 114.57 | 214.48 | 214.48 | 214.48 | 179.53 | |
| 7 | 198.60 | 198.60 | 198.60 | 106.96 | 210.03 | 210.03 | 240.03 | 180.45 | |
| 8 | 260.84 | 260.84 | 260.84 | 142.44 | 211.71 | 211.71 | 216.71 | 192.49 | |
| 9 | 342.54 | 342.54 | 342.54 | 125.33 | 213.75 | 213.75 | 217.75 | 188.52 | |
| 10 | 330.01 | 330.01 | 330.01 | 115.06 | 198.38 | 198.38 | 210.38 | 196.70 | |
| 11 | 239.27 | 239.27 | 239.27 | 134.57 | 214.48 | 214.48 | 214.48 | 210.53 | |
| 12 | 318.60 | 318.60 | 318.60 | 106.96 | 228.03 | 228.03 | 218.03 | 180.45 | |
| 13 | 360.84 | 360.84 | 360.84 | 122.44 | 222.71 | 222.71 | 222.71 | 192.49 | |
| 14 | 242.54 | 242.54 | 242.54 | 145.33 | 221.75 | 221.75 | 212.75 | 198.52 | |
| 15 | 250.01 | 250.01 | 250.01 | 135.06 | 213.38 | 213.38 | 213.38 | 200.70 | |
| 16 | 299.27 | 299.27 | 289.27 | 144.57 | 208.00 | 208.00 | 208.00 | 199.53 | |
| 17 | 318.60 | 318.60 | 318.60 | 106.96 | 246.03 | 246.03 | 206.03 | 198.45 | |
| 18 | 360.84 | 360.84 | 362.84 | 142.44 | 208.71 | 208.71 | 248.71 | 192.49 | |
| 19 | 242.54 | 242.54 | 242.54 | 125.33 | 234.75 | 234.75 | 244.75 | 189.52 | |
| 20 | 387.01 | 387.01 | 387.01 | 125.06 | 244.38 | 244.38 | 244.38 | 200.02 | |
| Mean± | 282.1± | 282.1± | 281.8± | 126.4± | 223.2± | 223.2± | 226.2± | 190.2± | |
| SD | 57.69 | 57.69 | 57.67 | 13.7*# | 18.84 | 18.84 | 18.53 | 9.986*# | |

Table 4b. Pre- and post- training lipid profile of obese female control and experimental subjects

Data were presented as mean ± standard deviation (SD); *= Statistical significance at p<0.05 when compared with pre-test of experimental subjects; #= Statistical significance at p<0.05 when compared with post-test of control subjects

| S/N | Burst (cm) | Chest (cm) | Abdomen (cm) | Waist (cm) | Hip (cm) | Thigh L (cm) | Thigh R (cm) | Calf L (cm) | Calf R (cm) | Upper Arm L | Upper Arm R | Upper Arm R | Lower Arm R |
|-------|---------------|---------------|-----------------|---------------|-------------|-----------------|-----------------|----------------|----------------|----------------|----------------|----------------|----------------|
| | | | | | | | | | | (cm) | (cm) | (cm) | (cm) |
| 1 | 121 | 109 | 112 | 118 | 114 | 67 | 69 | 43 | 43 | 41 | 40 | 30 | 30 |
| 2 | 99 | 90 | 90 | 108 | 111 | 57 | 58 | 37 | 38 | 30 | 29 | 26 | 25 |
| 3 | 118 | 102 | 106 | 118 | 125 | 59 | 62 | 45 | 43 | 37 | 37 | 29 | 30 |
| 4 | 99 | 92 | 85 | 107 | 111 | 62 | 64 | 39 | 39 | 34 | 34 | 28 | 27 |
| 5 | 88 | 84 | 73 | 94 | 89 | 53 | 53 | 33 | 33 | 26 | 27 | 25 | 25 |
| 6 | 78 | 90 | 84 | 115 | 119 | 65 | 65 | 41 | 41 | 31 | 32 | 28 | 28 |
| 7 | 91 | 87 | 83 | 111 | 113 | 66 | 66 | 41 | 41 | 33 | 34 | 22 | 24 |
| 8 | 120 | 110 | 112 | 116 | 117 | 70 | 68 | 47 | 47 | 37 | 38 | 19 | 19 |
| 9 | 92 | 89 | 79 | 105 | 108 | 57 | 62 | 38 | 38 | 30 | 30 | 22 | 25 |
| 10 | 110 | 100 | 98 | 120 | 120 | 69 | 68 | 36 | 36 | 39 | 38 | 18 | 18 |
| 11 | 110 | 100 | 98 | 120 | 120 | 69 | 68 | 36 | 36 | 39 | 38 | 18 | 18 |
| 12 | 92 | 89 | 79 | 105 | 108 | 57 | 62 | 38 | 38 | 30 | 30 | 22 | 25 |
| 13 | 120 | 110 | 112 | 116 | 117 | 70 | 68 | 47 | 47 | 37 | 38 | 19 | 19 |
| 14 | 91 | 87 | 83 | 111 | 113 | 66 | 66 | 41 | 41 | 33 | 34 | 21 | 24 |
| 15 | 78 | 90 | 84 | 115 | 119 | 65 | 65 | 41 | 41 | 31 | 32 | 28 | 28 |
| 16 | 88 | 84 | 73 | 94 | 89 | 53 | 53 | 33 | 33 | 26 | 27 | 25 | 25 |
| 17 | 99 | 92 | 85 | 107 | 111 | 62 | 64 | 39 | 39 | 34 | 34 | 28 | 27 |
| 18 | 118 | 102 | 106 | 118 | 125 | 59 | 62 | 45 | 43 | 37 | 37 | 29 | 30 |
| 19 | 99 | 90 | 90 | 108 | 111 | 57 | 58 | 37 | 38 | 30 | 29 | 26 | 25 |
| 20 | 121 | 109 | 112 | 118 | 114 | 67 | 69 | 43 | 43 | 41 | 40 | 30 | 30 |
| Mean± | 101.6± | 95.30± | 91.60± | 111.2± | 112.4± | 62.50± | 63.30± | 40.00± | 33.30± | 33.60± | 33.90± | 24.70± | 25.10± |
| SD | 14.56 | 9.027 | 13.84 | 7.743 | 9.282 | 5.624 | 5.583 | 4.155 | 5.059 | 4.500 | 3.959 | 4.131 | 3.959 |

Table 5a. Pre-test circumferential measurement of control subjects

| S/N | Burst (cm) | Chest (cm) | Abdomen (cm) | Waist (cm) | Hip (cm) | Thigh L (cm) | Thigh R (cm) | Calf L (cm) | Calf R (cm) | Upper Arm L (cm) | Upper Arm R (cm) | Upper Arm R (cm) | Lower Arm R (cm) |
|-----------|---------------|---------------|-----------------|---------------|-------------|-----------------|-----------------|----------------|----------------|------------------------|------------------------|------------------------|------------------------|
| 1 | 120 | 110 | 113 | 140 | 135 | 88 | 89 | 45 | 45 | 44 | 43 | 16 | 16 |
| 2 | 99 | 90 | 84 | 122 | 118 | 32 | 32 | 36 | 34 | 36 | 34 | 19 | 19 |
| 3 | 120 | 105 | 108 | 113 | 109 | 66 | 68 | 44 | 44 | 35 | 36 | 17 | 17 |
| 4 | 100 | 95 | 87 | 119 | 113 | 64 | 65 | 40 | 40 | 30 | 31 | 15 | 17 |
| 5 | 90 | 87 | 75 | 113 | 115 | 67 | 66 | 41 | 40 | 37 | 36 | 26 | 27 |
| 6 | 80 | 91 | 85 | 100 | 116 | 62 | 62 | 40 | 41 | 31 | 31 | 18 | 20 |
| 7 | 92 | 88 | 115 | 98 | 100 | 55 | 55 | 36 | 36 | 27 | 27 | 19 | 20 |
| 8 | 125 | 112 | 79 | 107 | 102 | 61 | 61 | 39 | 40 | 30 | 32 | 26 | 26 |
| 9 | 92 | 89 | 98 | 111 | 107 | 65 | 65 | 41 | 41 | 30 | 32 | 26 | 28 |
| 10 | 112 | 101 | 98 | 115 | 100 | 60 | 60 | 39 | 39 | 33 | 32 | 25 | 25 |
| 11 | 112 | 101 | 79 | 115 | 100 | 60 | 60 | 39 | 39 | 31 | 32 | 25 | 25 |
| 12 | 92 | 89 | 115 | 111 | 107 | 65 | 65 | 41 | 41 | 30 | 32 | 26 | 28 |
| 13 | 125 | 112 | 85 | 107 | 102 | 61 | 61 | 39 | 40 | 30 | 32 | 26 | 26 |
| 14 | 92 | 88 | 85 | 98 | 100 | 55 | 55 | 36 | 36 | 27 | 27 | 19 | 20 |
| 15 | 80 | 91 | 75 | 100 | 116 | 62 | 62 | 40 | 41 | 31 | 31 | 18 | 20 |
| 16 | 90 | 87 | 87 | 113 | 115 | 67 | 66 | 41 | 40 | 37 | 36 | 26 | 27 |
| 17 | 100 | 95 | 108 | 119 | 113 | 64 | 65 | 40 | 40 | 30 | 31 | 15 | 17 |
| 18 | 120 | 105 | 84 | 113 | 109 | 66 | 68 | 44 | 44 | 35 | 36 | 17 | 17 |
| 19 | 99 | 90 | 99 | 122 | 118 | 32 | 32 | 36 | 34 | 36 | 24 | 19 | 19 |
| 20 | 120 | 110 | 122 | 140 | 135 | 88 | 89 | 45 | 45 | 44 | 43 | 16 | 16 |
| Mean± | 103.0± | 96.80± | 92.90± | 112.5± | 113.6± | 63.00± | 63.30± | 40.80± | 33.00± | 34.00± | 34.20± | 24.60± | 25.10± |
| <u>2D</u> | 14.94 | 9.220 | 14.14 | 0.185 | 9.247 | 5.089 | 5.411 | 4.324 | 3.013 | 4.401 | 4.250 | 4.109 | 3.905 |

Table 5b. Post-test circumferential measurement of control subjects

| S/N | Burst | Chest (cm) | Abdomen | Waist | Hip (cm) | Thigh | Thigh B (cm) | Calf L | Calf R | Upper Arm I | Upper Arm R | Upper Arm B | Lower |
|-------|--------|---------------|---------|--------|-------------|--------|-----------------|--------|--------|----------------|----------------|----------------|--------|
| | (CIII) | (ciii) | (ciii) | (ciii) | (ciii) | | | (ciii) | (ciii) | (cm) | (cm) | (cm) | (cm) |
| 1 | 126 | 113 | 126 | 142 | 138 | 90 | 91 | 47 | 48 | 48 | 46 | 18 | 19 |
| 2 | 113 | 102 | 100 | 124 | 118 | 69 | 68 | 36 | 34 | 38 | 37 | 18 | 18 |
| 3 | 110 | 107 | 105 | 116 | 113 | 68 | 70 | 46 | 47 | 37 | 39 | 19 | 19 |
| 4 | 112 | 104 | 107 | 121 | 117 | 65 | 67 | 44 | 43 | 33 | 35 | 17 | 19 |
| 5 | 109 | 101 | 101 | 115 | 114 | 64 | 56 | 43 | 41 | 35 | 35 | 18 | 19 |
| 6 | 194 | 90 | 93 | 117 | 121 | 63 | 65 | 42 | 41 | 34 | 32 | 18 | 18 |
| 7 | 193 | 88 | 79 | 101 | 102 | 54 | 58 | 39 | 37 | 29 | 29 | 21 | 23 |
| 8 | 117 | 105 | 98 | 104 | 107 | 61 | 62 | 40 | 41 | 34 | 35 | 28 | 28 |
| 9 | 191 | 87 | 83 | 109 | 113 | 64 | 65 | 43 | 42 | 32 | 33 | 22 | 22 |
| 10 | 196 | 90 | 78 | 114 | 119 | 65 | 65 | 41 | 41 | 29 | 29 | 26 | 26 |
| 11 | 196 | 90 | 78 | 114 | 119 | 65 | 65 | 41 | 41 | 29 | 29 | 26 | 26 |
| 12 | 191 | 87 | 83 | 109 | 113 | 64 | 65 | 43 | 42 | 32 | 33 | 22 | 22 |
| 13 | 117 | 105 | 98 | 104 | 107 | 61 | 62 | 40 | 42 | 34 | 35 | 28 | 28 |
| 14 | 193 | 88 | 79 | 101 | 102 | 54 | 58 | 39 | 37 | 29 | 29 | 21 | 23 |
| 15 | 194 | 90 | 93 | 117 | 121 | 63 | 65 | 42 | 41 | 34 | 32 | 18 | 18 |
| 16 | 109 | 101 | 101 | 115 | 114 | 64 | 56 | 43 | 41 | 35 | 35 | 118 | 19 |
| 17 | 112 | 104 | 107 | 121 | 117 | 65 | 67 | 44 | 43 | 33 | 35 | 17 | 19 |
| 18 | 110 | 107 | 105 | 116 | 113 | 68 | 70 | 46 | 47 | 37 | 39 | 19 | 19 |
| 19 | 113 | 102 | 100 | 124 | 118 | 69 | 68 | 36 | 34 | 38 | 37 | 18 | 18 |
| 20 | 126 | 113 | 126 | 142 | 138 | 90 | 91 | 47 | 48 | 48 | 46 | 18 | 19 |
| Mean± | 106.1± | 98.70± | 97.00± | 116.3± | 116.2± | 66.30± | 66.7± | 42.10± | 41.50± | 34.90± | 35.00± | 20.70± | 21.50± |
| 30 | 06.11 | 0.945 | 14.20 | 11.18 | 9.30 | 9.039 | 9.314 | 3.194 | 4.007 | 5.320 | 4.077 | 3.0/8 | 2.014 |

Table 5c. Pre-test circumferential measurement of experimental subjects

| S/N | Burst (cm) | Chest (cm) | Abdomen (cm) | Waist (cm) | Hip (cm) | Thigh L (cm) | Thigh R (cm) | Calf L (cm) | Calf R (cm) | Upper Arm L (cm) | Upper Arm R (cm) | Upper Arm R (cm) | Lower Arm R (cm) |
|-------|---------------|---------------|-----------------|---------------|-------------|-----------------|-----------------|----------------|----------------|------------------------|------------------------|------------------------|------------------------|
| 1 | 120 | 110 | 122 | 140 | 135 | 88 | 89 | 45 | 45 | 44 | 43 | 16 | 16 |
| 2 | 110 | 100 | 99 | 122 | 118 | 32 | 32 | 36 | 34 | 36 | 34 | 19 | 19 |
| 3 | 100 | 105 | 100 | 113 | 109 | 66 | 68 | 44 | 44 | 35 | 36 | 17 | 17 |
| 4 | 110 | 99 | 102 | 119 | 113 | 64 | 65 | 40 | 40 | 30 | 31 | 15 | 17 |
| 5 | 103 | 86 | 97 | 113 | 115 | 67 | 66 | 41 | 40 | 37 | 36 | 26 | 27 |
| 6 | 89 | 85 | 84 | 100 | 116 | 62 | 62 | 40 | 41 | 31 | 31 | 18 | 20 |
| 7 | 90 | 100 | 78 | 98 | 100 | 55 | 55 | 36 | 36 | 27 | 27 | 19 | 20 |
| 8 | 115 | 110 | 95 | 107 | 102 | 61 | 61 | 39 | 40 | 30 | 32 | 26 | 26 |
| 9 | 87 | 82 | 81 | 111 | 107 | 65 | 65 | 41 | 41 | 30 | 32 | 26 | 28 |
| 10 | 93 | 87 | 84 | 115 | 100 | 60 | 60 | 39 | 39 | 33 | 32 | 25 | 25 |
| 11 | 93 | 87 | 84 | 115 | 100 | 60 | 60 | 39 | 39 | 31 | 32 | 25 | 25 |
| 12 | 87 | 82 | 81 | 111 | 107 | 65 | 65 | 41 | 41 | 30 | 32 | 26 | 28 |
| 13 | 115 | 110 | 95 | 107 | 102 | 61 | 61 | 39 | 40 | 30 | 32 | 26 | 26 |
| 14 | 90 | 100 | 78 | 98 | 100 | 55 | 55 | 36 | 36 | 27 | 27 | 19 | 20 |
| 15 | 89 | 85 | 84 | 100 | 116 | 62 | 62 | 40 | 41 | 31 | 31 | 18 | 20 |
| 16 | 103 | 86 | 97 | 113 | 115 | 67 | 66 | 41 | 40 | 37 | 36 | 26 | 27 |
| 17 | 110 | 99 | 102 | 119 | 113 | 64 | 65 | 40 | 40 | 30 | 31 | 15 | 17 |
| 18 | 100 | 105 | 100 | 113 | 109 | 66 | 68 | 45 | 44 | 35 | 36 | 17 | 17 |
| 19 | 110 | 100 | 99 | 122 | 118 | 32 | 32 | 36 | 34 | 36 | 24 | 19 | 19 |
| 20 | 120 | 110 | 122 | 140 | 135 | 88 | 89 | 45 | 45 | 44 | 43 | 16 | 16 |
| Mean± | 101.7± | 95.40± | 94.20± | 113.3± | 115.5± | 61.80± | 62.30± | 40.10± | 40.00± | 33.40± | 33.40± | 20.70± | 21.10± |
| SD | 11.15*# | 9.338* | 12.75*# | 11.44* | 10.31# | 13.49*# | 13.58* | 2.845* | 3.179# | 4.706 | 4.309* | 4.402# | 4.431# |

Table 5d. Post-test circumferential measurement of experimental subjects

Data were presented as mean ± standard deviation (SD); *= Statistical significance at p<0.05when compared with pre-test of experimental subjects; #= Statistical significance at p<0.05when compared with post-test of control subjects

| S/N | Biceps | Biceps | Triceps L | Triceps R | Chest | Abdomen(| Bi-iliac | Sub- | Inner | Inner | Calf L | Calf R |
|-------|--------|--------|-----------|-----------|--------|----------|----------|----------|---------|---------|--------|--------|
| | L (mm) | R (mm) | (mm) | (mm) | (mm) | mm) | (mm) | Scapular | Thigh L | Thigh R | (mm) | (mm) |
| | | | | | | | | (mm) | (mm) | (mm) | | |
| 1 | 35.7 | 40.0 | 32.4 | 29.8 | 27.6 | 32.9 | 35.5 | 21.1 | 18.8 | 15.3 | 17.4 | 18.4 |
| 2 | 35.9 | 41.9 | 26.6 | 21.0 | 35.5 | 25.3 | 30.1 | 25.8 | 18.4 | 16.7 | 13.2 | 14.6 |
| 3 | 31.8 | 32.3 | 18.7 | 20.1 | 31.5 | 19.0 | 24.9 | 18.9 | 18.6 | 16.0 | 19.6 | 16.5 |
| 4 | 26.5 | 27.5 | 23.1 | 23.0 | 32.3 | 29.9 | 32.3 | 28.7 | 20.2 | 19.9 | 13.4 | 13.3 |
| 5 | 45.2 | 35.1 | 38.2 | 27.8 | 29.6 | 41.3 | 38.4 | 25.0 | 30.9 | 31.0 | 12.6 | 16.3 |
| 6 | 33.1 | 37.1 | 22.7 | 26.3 | 34.0 | 37.1 | 26.5 | 23.1 | 18.2 | 14.6 | 11.0 | 11.8 |
| 7 | 34.2 | 22.9 | 18.6 | 25.6 | 25.0 | 23.9 | 34.5 | 30.7 | 32.0 | 28.1 | 18.9 | 19.6 |
| 8 | 40.5 | 36.7 | 19.0 | 17.0 | 27.2 | 32.5 | 30.2 | 21.0 | 18.8 | 22.0 | 15.8 | 19.0 |
| 9 | 28.0 | 30.5 | 25.1 | 20.0 | 26.4 | 26.0 | 27.0 | 24.7 | 17.1 | 16.4 | 18.6 | 16.6 |
| 10 | 32.4 | 29.0 | 26.3 | 26.1 | 27.0 | 27.8 | 30.0 | 27.5 | 20.5 | 22.0 | 13.5 | 12.1 |
| 11 | 32.4 | 29.0 | 26.3 | 26.1 | 27.0 | 27.8 | 30.0 | 27.5 | 20.5 | 22.0 | 13.5 | 12.1 |
| 12 | 28.0 | 30.5 | 25.1 | 20.1 | 26.4 | 26.0 | 27.0 | 24.7 | 17.1 | 16.4 | 18.6 | 16.6 |
| 13 | 40.5 | 36.7 | 19.0 | 17.0 | 27.2 | 32.5 | 30.2 | 21.0 | 18.8 | 22.0 | 15.8 | 19.0 |
| 14 | 34.2 | 22.9 | 18.6 | 25.6 | 25.0 | 23.9 | 34.5 | 30.7 | 32.0 | 28.1 | 18.9 | 19.6 |
| 15 | 33.1 | 37.1 | 22.7 | 26.3 | 34.0 | 37.1 | 26.5 | 23.1 | 18.2 | 14.6 | 11.0 | 11.8 |
| 16 | 45.2 | 35.1 | 38.2 | 27.8 | 29.6 | 41.3 | 38.4 | 25.0 | 30.9 | 31.0 | 12.6 | 16.3 |
| 17 | 26.5 | 27.5 | 23.1 | 23.0 | 32.3 | 29.9 | 32.3 | 28.7 | 20.2 | 19.9 | 13.4 | 13.3 |
| 18 | 31.8 | 32.3 | 18.7 | 21.0 | 31.5 | 19.0 | 24.9 | 18.9 | 18.6 | 16.0 | 19.6 | 16.5 |
| 19 | 35.9 | 41.9 | 26.6 | 21.0 | 35.5 | 25.3 | 30.1 | 25.8 | 18.4 | 16.7 | 13.2 | 14.6 |
| 20 | 35.7 | 40.0 | 32.4 | 29.8 | 27.6 | 32.9 | 35.5 | 21.1 | 18.8 | 15.3 | 17.4 | 18.4 |
| Mean± | 34.33± | 33.30± | 25.03± | 23.67± | 29.74± | 29.40± | 30.94± | 24.65± | 21.35± | 19.70± | 15.40± | 15.82± |
| SD | 5.367 | 5.779 | 6.101 | 3.966 | 3.429 | 6.452 | 4.165 | 3.165 | 5.271 | 5.169 | 2.973 | 2.718 |

Table 6a. Pre-test skin-fold measurement of control subjects

| S/N | Biceps | Biceps | Triceps L | Triceps | Chest | Abdomen(m | Bi-iliac | Sub- | Inner | Inner | Calf L | Calf R |
|------|----------|--------|-----------|---------|--------|-----------|----------|------------------|-----------------|-----------------|--------|--------|
| | L (mm) | R (mm) | (mm) | R (mm) | (mm) | m) | (mm) | Scapular (mm) | Thigh L (mm) | Thigh R (mm) | (mm) | (mm) |
| 1 | 30.0 | 26.9 | 22.7 | 19.2 | 28.9 | 19.8 | 30.8 | 23.1 | 18.2 | 20.4 | 15.8 | 17.0 |
| 2 | 35.0 | 40.7 | 25.0 | 22.4 | 40.1 | 27.0 | 34.5 | 24.4 | 18.9 | 18.0 | 14.2 | 15.8 |
| 3 | 32.5 | 40.1 | 20.7 | 30.5 | 37.8 | 25.1 | 30.4 | 27.9 | 18.5 | 16.8 | 21.5 | 19.0 |
| 4 | 27.4 | 30.2 | 25.0 | 27.0 | 40.2 | 32.9 | 37.4 | 31.2 | 21.9 | 20.0 | 15.6 | 15.5 |
| 5 | 38.4 | 39.3 | 40.1 | 32.8 | 30.6 | 39.3 | 40.3 | 27.1 | 30.9 | 32.7 | 15.4 | 18.2 |
| 6 | 32.3 | 39.1 | 30.1 | 28.3 | 41.7 | 39.1 | 28.8 | 27.3 | 20.2 | 18.2 | 12.1 | 12.8 |
| 7 | 34.5 | 27.0 | 20.6 | 27.0 | 30.5 | 29.3 | 37.5 | 32.1 | 30.2 | 28.3 | 19.6 | 20.4 |
| 8 | 41.0 | 39.9 | 22.0 | 24.5 | 27.2 | 32.5 | 30.2 | 27.0 | 18.9 | 22.1 | 19.0 | 20.9 |
| 9 | 30.5 | 36.7 | 27.1 | 25.2 | 34.4 | 28.9 | 30.0 | 28.7 | 20.0 | 22.6 | 21.6 | 18.6 |
| 10 | 35.4 | 32.0 | 26.8 | 26.3 | 30.7 | 28.7 | 40.1 | 29.5 | 22.5 | 23.0 | 17.5 | 12.1 |
| 11 | 35.4 | 32.0 | 26.8 | 26.3 | 30.7 | 28.7 | 40.1 | 29.5 | 22.5 | 23.0 | 17.5 | 12.1 |
| 12 | 30.5 | 36.7 | 27.1 | 25.2 | 34.4 | 28.9 | 30.0 | 28.7 | 20.0 | 22.6 | 21.6 | 18.6 |
| 13 | 41.0 | 39.9 | 22.0 | 24.5 | 27.2 | 32.5 | 30.2 | 27.0 | 18.9 | 22.1 | 19.0 | 20.9 |
| 14 | 34.5 | 27.0 | 20.6 | 27.0 | 30.4 | 29.3 | 37.5 | 32.1 | 30.2 | 28.3 | 19.6 | 20.4 |
| 15 | 32.3 | 39.1 | 30.1 | 28.3 | 41.7 | 39.1 | 28.8 | 27.3 | 20.2 | 18.2 | 12.1 | 12.8 |
| 16 | 38.4 | 39.3 | 40.1 | 32.8 | 30.6 | 39.3 | 40.3 | 27.1 | 30.9 | 32.7 | 15.4 | 18.2 |
| 17 | 27.4 | 30.2 | 25.0 | 27.0 | 40.2 | 32.9 | 37.4 | 31.2 | 21.9 | 20.0 | 15.6 | 15.5 |
| 18 | 32.5 | 40.1 | 20.7 | 30.5 | 37.8 | 25.1 | 30.4 | 27.9 | 18.5 | 16.8 | 21.5 | 19.0 |
| 19 | 35.0 | 40.7 | 25.0 | 22.4 | 40.1 | 27.0 | 34.5 | 24.4 | 18.9 | 18.0 | 14.2 | 15.8 |
| 20 | 30.0 | 26.9 | 22.7 | 19.2 | 28.9 | 19.8 | 30.8 | 23.1 | 18.2 | 20.4 | 15.8 | 17.0 |
| Mean | 34.33±5. | 33.30± | 25.03± | 23.67± | 29.74± | 29.40± | 30.94± | 24.65± | 21.35± | 19.70± | 15.40± | 15.82± |
| ±SD | 367 | 5.779 | 6.101 | 3.966 | 3.429 | 6.452 | 4.165 | 3.612 | 5.271 | 5.169 | 2.973 | 2.718 |

Table 6b. Post-test skin-fold measurement of control subjects

| S/N | Biceps L (mm) | Biceps R (mm) | Triceps L (mm) | Triceps R (mm) | Chest (mm) | Abdomen (mm) | Bi-iliac (mm) | Sub- Scapular | Inner Thigh L | Inner Thigh R | Calf L (mm) | Calf R (mm) |
|-------|------------------|------------------|-------------------|-------------------|---------------|-----------------|------------------|------------------|------------------|------------------|----------------|----------------|
| | | | | | | | | (mm) | (mm) | (mm) | | |
| 1 | 37.0 | 23.8 | 20.1 | 17.2 | 27.2 | 26.07 | 37.6 | 17.0 | 14.5 | 22.7 | 20.3 | 17.0 |
| 2 | 28.0 | 26.9 | 18.1 | 19.2 | 42.9 | 25.8 | 30.8 | 20.5 | 23.3 | 20.3 | 19.5 | 19.5 |
| 3 | 37.8 | 25.4 | 20.0 | 19.0 | 27.0 | 30.0 | 28.2 | 20.0 | 15.2 | 15.2 | 15.0 | 17.8 |
| 4 | 20.5 | 20.0 | 23.2 | 20.3 | 23.0 | 30.8 | 34.5 | 20.7 | 18.0 | 18.0 | 17.1 | 17.8 |
| 5 | 32.3 | 31.6 | 26.4 | 25.6 | 28.4 | 34.8 | 28.4 | 20.7 | 16.8 | 16.8 | 16.5 | 13.2 |
| 6 | 36.4 | 31.1 | 21.9 | 24.4 | 36.9 | 35.5 | 32.5 | 20.6 | 22.1 | 22.1 | 11.4 | 13.6 |
| 7 | 30.0 | 29.4 | 22.4 | 20.9 | 24.0 | 20.7 | 25.5 | 22.9 | 14.7 | 14.7 | 15.8 | 15.6 |
| 8 | 20.9 | 20.6 | 29.8 | 17.0 | 28.9 | 19.8 | 20.0 | 17.9 | 19.9 | 19.9 | 18.0 | 20.4 |
| 9 | 34.2 | 22.9 | 18.6 | 26.6 | 35.1 | 35.9 | 28.5 | 20.7 | 32.0 | 32.0 | 10.4 | 12.8 |
| 10 | 33.1 | 37.1 | 22.7 | 26.3 | 34.0 | 37.1 | 34.8 | 23.1 | 18.2 | 18.2 | 10.6 | 16.6 |
| 11 | 33.1 | 37.1 | 22.7 | 26.3 | 34.0 | 37.1 | 34.8 | 23.1 | 18.2 | 18.2 | 10.6 | 16.6 |
| 12 | 34.2 | 22.9 | 18.6 | 17.0 | 35.1 | 35.9 | 28.5 | 20.7 | 32.0 | 32.0 | 10.4 | 12.8 |
| 13 | 20.9 | 20.6 | 29.9 | 20.9 | 28.9 | 19.8 | 20.0 | 17.9 | 19.9 | 19.9 | 18.0 | 20.4 |
| 14 | 30.0 | 29.4 | 22.4 | 24.4 | 24.0 | 20.7 | 25.5 | 22.9 | 14.7 | 14.7 | 15.8 | 15.6 |
| 15 | 36.4 | 31.1 | 21.9 | 25.6 | 36.9 | 35.5 | 32.5 | 20.6 | 22.1 | 22.1 | 11.4 | 13.6 |
| 16 | 32.3 | 31.6 | 26.4 | 20.3 | 28.4 | 34.8 | 28.4 | 20.7 | 16.8 | 16.8 | 16.5 | 13.2 |
| 17 | 20.5 | 20.0 | 23.2 | 19.0 | 23.0 | 30.8 | 34.5 | 20.7 | 18.0 | 18.0 | 17.1 | 17.8 |
| 18 | 37.8 | 25.4 | 20.0 | 19.2 | 27.0 | 30.0 | 28.2 | 20.0 | 15.2 | 15.2 | 15.0 | 17.8 |
| 19 | 28.0 | 26.9 | 18.1 | 17.2 | 42.9 | 25.8 | 30.8 | 20.5 | 23.3 | 20.3 | 19.5 | 19.5 |
| 20 | 37.0 | 23.8 | 20.1 | 118 | 27.2 | 19.9 | 37.6 | 17.0 | 14.5 | 22.7 | 20.3 | 17.0 |
| Mean± | 31.02± | 26.68± | 22.32± | 21.65± | 30.74± | 35.2± | 30.08± | 24.41± | 19.43± | 20.78± | 18.67± | 16.81± |
| SD | 6.070 | 5.283 | 3.489 | 3.644 | 6.123 | 6.805 | 4.986 | 1.830 | 5.327 | 4.266 | 3.590 | 2.616 |

Table 6c. Pre-test skin-fold measurement of experimental subjects

| S/N | Biceps L | Biceps | Triceps | Triceps | Chest | Abdomen(| Bi-iliac | Sub- | Inner | Inner | Calf L | Calf R |
|-------|----------|--------|---------|---------|--------|----------|----------|----------|---------|------------------|--------|--------|
| | (mm) | R (mm) | L (mm) | R (mm) | (mm) | mm) | (mm) | Scapular | | I nign B (mm) | (mm) | (mm) |
| 4 | | | | 40.5 | | 01.0 | | (1111) | (11111) | | | 40.0 |
| 1 | 37.2 | 23.8 | 20.5 | 18.5 | 29.5 | 21.6 | 39.5 | 18.1 | 16.5 | 22.7 | 22.5 | 18.2 |
| 2 | 32.4 | 27.8 | 26.3 | 26.1 | 26.8 | 26.8 | 35.5 | 23.3 | 18.4 | 17.4 | 23.7 | 17.1 |
| 3 | 26.7 | 40.7 | 19.4 | 14.0 | 27.1 | 32.9 | 20.2 | 21.1 | 15.8 | 15.3 | 15.1 | 19.2 |
| 4 | 18.1 | 23.0 | 25.3 | 20.5 | 25.0 | 24.1 | 36.1 | 34.3 | 20.0 | 22.6 | 18.6 | 14.1 |
| 5 | 27.5 | 26.5 | 21.8 | 19.8 | 27.9 | 33.6 | 28.4 | 28.8 | 18.8 | 20.0 | 20.3 | 21.1 |
| 6 | 28.7 | 27.5 | 18.4 | 14.6 | 27.5 | 29.2 | 24.2 | 21.9 | 14.9 | 12.9 | 12.2 | 12.0 |
| 7 | 27.0 | 30.3 | 24.9 | 20.0 | 24.4 | 24.1 | 25.2 | 23.7 | 15.8 | 15.3 | 18.6 | 16.6 |
| 8 | 20.9 | 22.6 | 17.3 | 17.5 | 30.2 | 19.8 | 22.5 | 17.9 | 21.3 | 21.5 | 21.3 | 18.4 |
| 9 | 21.4 | 20.8 | 15.9 | 24.5 | 25.0 | 23.6 | 34.5 | 32.9 | 29.0 | 21.3 | 18.9 | 19.6 |
| 10 | 26.1 | 17.8 | 17.4 | 15.5 | 26.9 | 18.0 | 26.5 | 22.2 | 10.7 | 14.6 | 11.0 | 11.8 |
| 11 | 26.1 | 17.8 | 17.4 | 15.6 | 26.9 | 18.0 | 26.5 | 22.2 | 10.7 | 14.6 | 11.0 | 11.8 |
| 12 | 21.4 | 20.8 | 15.9 | 24.5 | 25.0 | 23.6 | 34.5 | 32.9 | 29.0 | 21.3 | 18.9 | 19.6 |
| 13 | 20.9 | 22.6 | 17.3 | 17.5 | 30.2 | 19.8 | 22.5 | 17.9 | 21.3 | 21.5 | 21.3 | 18.4 |
| 14 | 27.0 | 30.3 | 24.9 | 20.0 | 24.4 | 24.1 | 25.2 | 23.7 | 15.8 | 15.3 | 18.6 | 16.6 |
| 15 | 28.7 | 27.5 | 18.4 | 14.6 | 27.5 | 29.2 | 24.2 | 21.9 | 14.9 | 12.9 | 12.2 | 12.0 |
| 16 | 27.5 | 26.5 | 21.8 | 19.8 | 27.9 | 33.6 | 28.4 | 28.8 | 18.8 | 20.0 | 20.3 | 21.1 |
| 17 | 18.1 | 23.0 | 25.3 | 20.5 | 25.0 | 24.1 | 36.1 | 34.3 | 20.0 | 22.6 | 18.6 | 14.1 |
| 18 | 26.7 | 40.7 | 19.4 | 14.0 | 27.1 | 32.9 | 20.2 | 21.1 | 15.8 | 15.3 | 15.1 | 19.2 |
| 19 | 32.4 | 27.8 | 26.3 | 26.1 | 26.8 | 20.63s | 35.5 | 23.3 | 18.4 | 17.4 | 23.7 | 17.1 |
| 20 | 37.2 | 23.8 | 20.5 | 18.5 | 29.5 | 21.6 | 39.5 | 18.1 | 16.5 | 22.7 | 22.5 | 18.2 |
| Mean± | 26.60± | 26.08± | 20.72± | 19.25± | 27.25± | 29.03± | 29.26± | 20.41± | 18.12± | 18.36± | 15.4± | 16.43± |
| SD | 5.462*# | 6.155# | 3.611*# | 3.767*# | 3.52*# | 4.471* | 6.458 | 5.543*# | 4.717# | 3.578* | 3.96* | 2.61 |

Table 6d. Post-test skin-fold measurement of experimental subjects

Data were presented as mean ± standard deviation (SD); *= Statistical significance at p<0.05when compared with pre-test of experimental subjects; #= Statistical significance at p<0.05when compared with post-test of control subjects

This study revealed that there was an increase in blood alucose in both control and experimental subjects during pre-test period. This could be as a result of excess glucose being stored in the skeletal muscle and the liver, this process is mediated by the hormone insulin. Exercise enhances insulin sensitivity by activating 5' AMPactivated protein kinase (AMPK) for glucose absorption (Mann et al., 2014). During aerobic exercise, the glucose stored in the muscle and the liver in form of alycogen is broken down to as energy in form of adenosine serve triphosphate (ATP) to fuel the physical exercise (Alghannam et al., 2021). The breakdown of glycogen into glucose in the skeletal muscle (gluconeogenesis) and liver is regulated by the hormone insulin and glucagon.

This study reveals that blood glucose was lowered with aerobic exercise in female obese experimental subjects than the female obese control subjects. Previous studies showed that aerobic exercise is an effective strategy for improvina glucose metabolism outcome (Poehlman et al 2000, Rahimi et al., 2020). Also previous study reported that aerobic exercise improves muscle capillary which helps glucose to diffuse from capillary into muscle cells, changes the muscle fibre type, favouring substrate oxidation and increases the quantity and size of mitochondria (Haas and Nwadozi, 2015). Aerobic exercise increases the expression and activity of key signaling protein involved in the regulation of glucose uptake and metabolism in skeletal muscle(such as GLUT4 translocation mediated by AMPK) and increases lipid oxidation (Hawley et al., 2008).

This study revealed that aerobic exercise decrease lipid profile (triglyceride, low-density lipo-protein, and total cholesterol and increase high density lipoprotein which is a good cholesterol in the experimental subjects but the reverse is the case in obese control subjects who do not participate in the aerobic exercise. Previous evidence shows that exercise have been found to positively affect lipid metabolism by improving lipid profile parameters such as triglyceride, low density lipo- protein, high density lipo-protein, and total cholesterol this is in line with study carried out by (Gordon et al., 2014).Aerobic exercise has been shown to enhance insulin sensitivity in obese individual improved insulin sensitivity helps to regulate blood glucose levels and reduces the production of triglyceride and very low density lipo-protein in the liver ,leading to favorable changes in the lipid

profile (Bird and Hawley, 2017). Aerobic exercise can potentially reduce visceral fat stored around the organs in the abdominal area. visceral fats is metabolically active and contributes to increased triglyceride and low density lipo-protein levels often called bad cholesterol, exercise can improve the lipid profile by reducing visceral fat and decreasing triglyceride and low density lipoprotein (Neeland et al., 2024). Regular exercise including aerobic exercise has increased the level high density lipo-protein level, commonly known as good cholesterol. High density lipoprotein help in transport excess cholesterol from the peripheral tissues to the liver for excretion, thus reducing the risk of cardio vascular diseases. Exercise can modify the size and density of low level density lipo- protein particles making them less atherogenic that less likely to cause plague build up in the arteries (Franczyk et al., 2023). Following Aerobic exercise the mechanism responsible for the acute rise in high density lipo-protein is likely associated with the catabolism of lipo-proteins rich triglyceride through lipo-protein lipase LPL (Ahotupa, 2024). Increase activity of lipo-protein lipase following exercise results in hydrolysis of chylomicron and very low density lipo-protein, thereby decreasing plasma triglyceride (Calder and Yaqoob, 2023), subsequently surface remnants and are converted into nascent high density lipo-protein, following the exercise the levels of high density lipo-protein can increase. Aerobic exercises primarily utilize fat as fuel for exercise by increasing the intensity and duration of exercise, patients can enhance fat oxidation, reducing overall body fat and improving their lipid profile (Talanian et al., 2007).

5. CONCLUSION

Aerobic exercise reduced body weight, lipid profile and blood glucose levels by making use of stored glucose and fats in the body as source of energy for the physical activities thereby reducing the risk factors associated with obesity, cardiovascular diseases and lots more.

This study confirms the data in the literature on the adverse effect of sedentary lifestyle on the body system.

Also, it revealed the significance of aerobic exercise in reducing body weight and risk of obesity, cardiovascular diseases, hypertension, stroke and lots more. Exercise, if it can be process into drugs informs of capsules, syrup and injection it could be used as therapeutic agent to prevent and cure several many diseases.

Future studies could consider more diverse populations, other types of aerobic exercise and qualitative insights for better establishment of the impact of physical activities on obesity management.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative Al technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

CONSENT

As per international standards or university standards, Participants' written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

Ethical Approval was obtained from the Ministry of Health Agodi, Ibadan, Oyo state, with Ethical Research Approval Number: NHREC/OYOSHRIEC/10/11/22

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Ahotupa, M. (2024). Lipid oxidation products and the risk of cardiovascular diseases: Role of lipoprotein transport. *Antioxidants, 13*(5), 512.
- Alghannam, A. F., Ghaith, M. M., & Alhussain, M.
 H. (2021). Regulation of energy substrate metabolism in endurance exercise. *International Journal of Environmental Research and Public Health, 18*(9), 4963.
- Archer, E., Shook, R. P., Thomas, D. M., Church, T. S., Katzmarzyk, P. T., & Hebert, J. R. (2013). 45-year trends in women's use of time and household management energy expenditure. *PLoS One*, *8*, e56620.
- Bird, S. R., & Hawley, J. A. (2017). Update on the effects of physical activity on insulin sensitivity in humans. *BMJ Open Sport & Exercise Medicine*, 2(1), e000143.

- Blair, S. N., Archer, E., & Hand, G. A. (2013). Commentary: Luke and Cooper are wrong: Physical activity has a crucial role in weight management and determinants of obesity. *International Journal of Epidemiology, 42*, 1836–1838.
- Calder, P. C., & Yaqoob, P. (2023). 8 Fat metabolism. *Human Nutrition, 161*.
- Chiu, C. H., Ko, M. C., Wu, L. S., Yeh, D. P., Kan, N. W., Lee, P. F., & Ho, C. C. (2017). Benefits of different intensity of aerobic exercise in modulating body composition among obese young adults: A pilot randomized controlled trial. *Health and Quality of Life Outcomes, 15*, 1-9.
- Church, T. S., LaMonte, M. J., Barlow, C. E., & Blair, S. N. (2005). Cardiorespiratory fitness and body mass index as predictors of cardiovascular disease mortality among men with diabetes. *Archives of Internal Medicine, 165*, 2114–2120.
- Fenkçi, S., Sarsan, A., Rota, S., & Ardıç, F. (2006). Effects of resistance or aerobic exercises on metabolic parameters in obese women who are not on a diet. *Advances in Therapy, 23*, 404-413.
- Fogelholm, M., Mannisto, S., Vartiainen, E., & Pietinen, P. (1996). Determinants of energy balance and overweight in Finland 1982 and 1992. *International Journal of Obesity and Related Metabolic Disorders*, 20, 1097–1104.
- Franczyk, B., Gluba-Brzózka, A., Ciałkowska-Rysz, A., Ławiński, J., & Rysz, J. (2023). The impact of aerobic exercise on HDL quantity and quality: A narrative review. *International Journal of Molecular Sciences, 24*(5), 4653.
- Gordon, B., Chen, S., & Durstine, J. L. (2014). The effects of exercise training on the traditional lipid profile and beyond. *Current Sports Medicine Reports, 13*(4), 253-259.
- Haas, T. L., & Nwadozi, E. (2015). Regulation of skeletal muscle capillary growth in exercise and disease. *Applied Physiology*, *Nutrition, and Metabolism, 40*(12), 1221-1232.
- Hawley, J. A., & Lessard, S. J. (2008). Exercise training–induced improvements in insulin action. *Acta Physiologica (Oxford), 192*(1), 127-135.
- Hill, J. O., & Peters, J. C. (2013). Commentary: Physical activity and weight control. International Journal of Epidemiology, 42, 1840–1842.
- Hojjat, T. A., & Hojjat, T. A. (2021). Introduction: The magnitude of the obesity problem. *The*

Economics of Obesity: Poverty, Income Inequality, and Health, 1-9.

- Karvonen, M. J. (1957). The effects of training on heart rate: A longitudinal study. *Annales Medicinae Experimentalis et Biologiae Fenniae*, *35*(3), 307–315.
- Lee, M., Wu, Y., & Fried, S. (2013). Adipose tissue heterogeneity: Implication of depot differences in adipose tissue for obesity complications. *Molecular Aspects of Medicine*, *34*(1), 1-11.
- Li, H.-L., Wu, X., Xu, A., & Hoo, R. L.-C. (2021). A-FABP in metabolic diseases and the therapeutic implications: An update. *International Journal of Molecular Sciences, 22*, 9386.
- Luke, A., & Cooper, R. S. (2013). Physical activity does not influence obesity risk: Time to clarify the public health message. *International Journal of Epidemiology, 42*, 1831–1836.
- Mann, S., Beedie, C., Balducci, S., Zanuso, S., Allgrove, J., Bertiato, F., et al. (2014). Changes in insulin sensitivity in response to different modalities of exercise: A review of the evidence. *Diabetes Metabolism Research Reviews, 30*, 257–268.
- Neeland, I. J., Lim, S., Tchernof, A., Gastaldelli, A., Rangaswami, J., Ndumele, C. E., & Després, J. P. (2024). Metabolic syndrome. *Nature Reviews Disease Primers, 10*(1), 77.
- Ng, M., Fleming, T., Robinson, M., Thomson, B., Graetz, N., & Margono, C. (2014). Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: A systematic analysis for the Global Burden of Disease Study 2013. *Lancet, 384*, 766–781.
- Ng, S. W., & Popkin, B. M. (2012). Time use and physical activity: A shift away from movement across the globe. *Obesity Reviews, 13*, 659–680.

- Plowman, S. A., & Smith, D. L. (2007). *Exercise* physiology for health, fitness and performance (2nd ed.).
- Poehlman, E. T., Dvorack, R. V., Dervino, W. F., Brochu, M., & Ades, P. A. (2000). Effect of exercise, resistance training, and endurance training on insulin sensitivity in non-obese, young women: A controlled randomized trial. *Endocrinology and Metabolism, 85*(7), 2463-2468.
- Prentice, A. M. (2006). The emerging epidemic of obesity in developing countries. *International Journal of Epidemiology, 35*, 93–99.
- Rahim, G. R. M., Bijeh, N., & Rashidalmir, A. (2020). Effect of exercise training on serum peptin, undercarboxylated osteocalcin, and high molecular weight adiponectin in adults with metabolic syndrome. *Experimental Physiology*, 105(3), 449-459.
- Stasiulis, A., Mockiene, A., Vizbaraite, O., & Mockus, P. (2004). Aerobic exercise induced changes in body composition and blood lipids in young women. *Medicina (Kaunas), 46*, 129-134.
- Talanian, J. L., Galloway, S. D., Heigenhauser, G. J., Bonen, A., & Spriet, L. L. (2007).
 Two weeks high intensity aerobic interval training increases the capacity for fat oxidation during exercise in women. *Journal of Applied Physiology*.
- Thorp, A. A., Owen, N., Neuhaus, M., & Dunstan, D. W. (2011). Sedentary behaviors and subsequent health outcomes in adults: A systematic review of longitudinal studies, 1996–2011. American Journal of Preventive Medicine, 41, 207–215.
- World Health Organization. (2016). *Physical activity strategy for the WHO European Region 2016-2025.*

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/129229