



Plasma Lipid Lowering Potential of Carrot (*Daucus carota*) Extract in Male Wistar Rats

Onyebuchi Obia ^{a*}, Regina Oripirisiki Kalio ^b,
Popnen Godwin Porobe Tee ^a and Godspower C. Onyeso ^a

^a Department of Human Physiology, faculty of Basic Medical Sciences, College of Medical Sciences, Rivers State University, Nkpolu-Oroworukwo, Port Harcourt, Nigeria.

^b Department of Human Physiology, faculty of Basic Medical Sciences, College of Health Sciences, University of Port Harcourt, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajrimps/2025/v14i1291>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/129295>

Original Research Article

Received: 08/11/2024
Accepted: 10/01/2025
Published: 14/01/2025

ABSTRACT

Carrots (*Daucus carota*) are widely consumed for their nutritional and medicinal benefits. The aim of the present study was to investigate the effects of *Daucus carota* on lipid profile. The study involved 28 male wistar rats separated into 4 groups of 7 rats each. Group 1 served as control and was given distilled water, whilst groups 2, 3 and 4 served as test groups and were given aqueous extract of *Daucus carota* at daily doses of 200mg/kg, 400mg/kg and 600mg/kg respectively. The experiment lasted for 28 days and thereafter, blood samples collected for determination of lipid profile [total cholesterol (TC), triglyceride (TG), high density lipoprotein (HDL) and low density

*Corresponding author: E-mail: onyebuchi.obia@ust.edu.ng;

Cite as: Obia, Onyebuchi, Regina Oripirisiki Kalio, Popnen Godwin Porobe Tee, and Godspower C. Onyeso. 2025. "Plasma Lipid Lowering Potential of Carrot (*Daucus Carota*) Extract in Male Wistar Rats". *Asian Journal of Research in Medical and Pharmaceutical Sciences* 14 (1):18-23. <https://doi.org/10.9734/ajrimps/2025/v14i1291>.

lipoprotein (LDL)] using standard laboratory techniques. Results showed that oral administration of all three doses of carrot extracts significantly reduced the plasma concentrations of both TC and TG but caused no significant changes in the LDL levels. However, the higher doses; 400mg/kg and 600mg/kg significantly increased the HDL levels in wistar rats. The lipid lowering potential of *Daucus carota* could be based on its fibre content as well as its antioxidant content which possibly enhanced fecal excretion of lipids. Conclusively, moderate consumption of *Daucus carota* decreased the plasma levels of TC and TG, increased HDL but had no significant effect on the LDL.

Keywords: Lipid lowering; potential; daucus carota; wistar rats.

1. INTRODUCTION

Carrot is an essential and popular root vegetable which was first used as medicine and later gradually used as food (Carlos and Dias, 2014; Bahrami et al., 2018). There are different varieties of carrots depending on the colour of their roots; orange, white, black, yellow, purple and red (Que et al., 2019; Purkiewicz et al., 2020), with predominantly more of the orange-coloured variety. It is a good source of some bioactive compounds which are beneficial to health including dietary fibre and micronutrients such as vitamins, minerals and antioxidants (Tanaka et al., 2012; Augpole et al., 2014; Fiedor and Burda, 2014; Bystrická et al., 2015). The colour of the root significantly affects the presence of bioactive compounds. For instance, the orange carrot root contains increased concentration of α -carotene and also very rich source of β -carotene, the black carrot root is a rich source of anthocyanins, red carrot root is endowed with lycopene whereas the yellow variety has been shown to accumulate lutein (Alasalvar et al., 2001; Molldrem et al., 2004; Stange and Rodriguez-Concepcion, 2015). Furthermore, the growing and season conditions, the degree of ripeness of carrots as well as part of the root to some extent also influence the presence of bioactive compounds (Molldrem et al., 2004). Carrots are generally rich in carotenoids (Hager and Howard, 2006) as well as phenols (Sharma et al., 2012; Leja et al., 2013; Ismail et al., 2023). Most fruits and vegetables are endowed with abundant phytochemicals, pro-vitamins and vitamins which protects against cellular damage and thus prevent certain diseases (Myojin et al., 2008; Phan et al., 2018; Zhu et al., 2018).

Plasma cholesterol is derived from both hepatic synthesis and diet, although the body balances the absorption of cholesterol by reducing its synthesis (Lecerf and Lorgeril, 2011). The main dietary sources of cholesterol include meat, eggs

and milk. However, there may be racial, gender and age variations in the average consumption of cholesterol (Xu et al., 2018). Studies have linked excessive dietary cholesterol with higher risk of cardiovascular disease (Wilde et al., 2000; Ingelsson et al., 2007; Duan et al., 2018; Carson et al., 2020; Wali et al., 2020; Guo et al., 2022; Stellaard, 2022; Zhao et al., 2022). Nutritional interventions and lifestyle changes have over the years been used in the prevention and management of these cardiovascular diseases. These interventions include, enriching the diet with dietary fibres, plant proteins and unsaturated fats (Kirkpatrick et al., 2023), aside increasing physical activity to prevent obesity. Healthy dietary patterns including the popularized Dietary Approaches to Stop Hypertension (DASH) which are relatively low in cholesterol should be of utmost priority. These patterns involve the consumption of vegetables, fruits, nuts, whole grains, seeds, low-fat or fat-free dairy products, lean protein sources and liquid vegetable oils (Carson et al., 2020). Hence, the dietary patterns approach would improve diet quality and ultimately promote cardiovascular health.

The aim of the present study was to determine the effect of carrot juice on the plasma lipid profile of wistar rats.

2. MATERIALS AND METHODS

The experiment was carried out at animal house of the department of Human Physiology, faculty of Basic Medical Sciences, University of Port Harcourt in the year 2019. A total of twenty eight male wistar rats weighing 120 to 150g were purchased and acclimatized for a period of two weeks. These animals were grouped and housed in plastic cages and allowed to feed and drink *ad libitum* with Top feed Finisher mash and clean water. Their immediate environment (beddings) was changed daily, the temperature of the environment kept at normal conditions while the external environment was cleaned and disinfected regularly.

2.1 Preparation and Administration of Carrot Extract

Mature carrot tubers were bought from Oil Mill market in Obio Akpor Local Government Area of Rivers State, Nigeria. The plant was identified at the department of Plant Science and Biotechnology, University of Port Harcourt; *Daucus carota* L, in the family; Apiceace with assigned herbarium number; UPH/C/132. The tubers were washed with water to remove soil particles. About 2kg of the fresh carrot was cut into tiny pieces and air dried for seven days. The dried carrots were blended using a blender and carefully poured into a maceration jar containing four liters of water. The mixture was allowed to macerate for 24hours after which a Whatman filter (20-25µm, pore size) was used to get a clear filtrate. The filtrate was now poured into an evaporating dish and dried in a water bath at 45°C to obtain a semi-solid aqueous extract of *Daucus carota*. The dosages administered in the study were based on the lethal dose (LD50) of 5000mg/kg which was previously determined (Ayeni et al., 2019). Following acclimatization, the wistar rats were weighed and separated into four groups of seven rats each. Group 1 served as control and was given distilled water, whilst groups 2, 3 and 4 served as test groups and were given aqueous extract of *Daucus carota* at daily doses of 200mg/kg, 400mg/kg and 600mg/kg respectively. The experiment lasted for 28 days. Thereafter the animals were sacrificed under anesthesia and blood samples collected via cardiac puncture for determination of lipid profile [total cholesterol (TC), triglyceride (TG), high density lipoprotein (HDL) and low density lipoprotein (LDL)] using spectrophotometric methods with appropriate reagents.

Statistical analysis was done using SPSS software version 22.0. Results were presented in Tables. Continuous variables were expressed as mean ± Standard error of mean (SEM). Statistical difference was determined using analysis of variance (ANOVA) and at $p < 0.05$.

3. RESULTS AND DISCUSSION

The results of our study showed that the three concentrations of carrot extract respectively caused significant reduction in TC and TG levels compared to the control. Carrots and other plant products (including fruits, vegetables, nuts, seeds and grains) contain dietary fibers (Anne et al., 2006) which are known to decrease the absorption of cholesterol from the intestines (Nicolle et al., 2003; Feingold, 2024; Obia and Eifuobhokhan, 2024). The reduction in TC and TG in the present study could be attributed to the presence of dietary fibers with reduced absorption of cholesterol and possibly increasing the excretion of cholesterol in faeces (Obia and Eifuobhokhan, 2024; Eifuobhokhan et al., 2024). Previous studies suggest that carotenoids and other antioxidants present in carrot and other plant products improves vascular function, reduces plasma lipids and prevents oxidative cellular damage and thus greatly reduces the risk of cardiovascular disease (Obia and Eifuobhokhan, 2024; Cheng et al., 2017; Obia and Asuquo, 2018; Obia et al., 2018; Aune, 2019; Chinko et al., 2023) and prevent premature deaths. Triglyceride lowering has been linked to reduction in cardiovascular disease risk (Marston et al., 2019).

The 200mg/kg of the extract had no significant effect on the plasma HDL levels but higher doses; 400mg/kg and 600mg/kg of the extract respectively increased the HDL levels in a dose-dependent manner. A lowering of the LDL with concomitant increase in the HDL is an important factor in the risk assessment of cardiovascular disease (Sirtori and Fumagalli, 2006). HDL is known to be the 'good' cholesterol and whatever factor that increases its plasma concentration will potentially lower the risk of cardiovascular disease (Kjeldsen et al., 2021). The result from present study therefore suggests that moderate consumption of carrot extract would potentially minimize the risk of cardiovascular disease. However, all three concentrations of the extract caused no significant changes in the plasma levels of LDL compared to control even though slight reduction in LDL was observed as dosage of the extract increased.

Table 1. Effect of Carrot extract on plasma lipid profile of wistar rats

Group	TC (mmol/l)	TG (mmol/l)	HDL (mmol/l)	LDL (mmol/l)
Control	5.49±0.23	1.57±0.05	0.80±0.07	2.13±0.14
200mg/kg	4.20±0.33*	1.21±0.07*	0.97±0.09	2.12±0.27
400mg/kg	3.30±0.26*	1.08±0.14*	1.26±0.11*	1.76±0.23
600mg/kg	3.28±0.37*	0.93±0.05*	1.67±0.07*	1.78±0.15

* Significantly different compared to control ($p < 0.05$)

4. CONCLUSION

Conclusively, oral administration of all three doses of carrot extracts for 28 days significantly reduced the plasma concentrations of both total cholesterol and triglyceride but caused no significant changes in the LDL levels. However, the higher doses; 400mg/kg and 600mg/kg significantly increased the HDL levels in wistar rats.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

CONSENT

It is not applicable.

ETHICAL APPROVAL

Ethical approval was obtained from the university of port harcourt research ethics committee with approval number; uph/ceremad/rec/mm67/012.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Carlos, J., Dias, S. (2014). Nutritional and Health Benefits of Carrots and Their Seed Extracts. *Food and Nutrition Sciences*, 5: 2147-2156.
- Bahrami, R., Ghobadi, A., Behnoud, N., & Akhtari, E. (2018). Medicinal properties of *Daucus carota* in traditional Persian medicine and modern phytotherapy. *Journal of Biochemical Technology*. Special issue (2), 107-114.
- Que, F., Hou, X. L., Wang, G. L., Xu, Z. S., Tan, G. F., Li, T., Wang, Y. H., Khadr, A., & Xiong, A. S. (2019). Advances in research on the carrot, an important root vegetable in the Apiaceae family. *Horticulture research*, 6, 69.
- Purkiewicz, A., Ciborska, J., Tanska, M., Narwojsz, A., Starowicz, M., Przybyłowicz, K. E., & Sawicki, T. (2020). The Impact of the Method Extraction and Different Carrot

- Variety on the Carotenoid Profile, Total Phenolic Content and Antioxidant Properties of Juices. *Plants*, 9, 1759.
- Tanaka, T., Shnimizu, M., & Moriwaki, H. (2012). Cancer chemoprevention by carotenoids. *Molecules* (Basel, Switzerland), 17 (3), 3202–3242.
- Augpole, I., Rackejeva, T., Kruma, Z., Dimins, F. (2014). Shredded carrots quality providing by treatment with Hydrogen peroxide. 9th Baltic Conference on “Food for Consumer Well-Being” *FOODBALT 2014*, 150-154.
- Fiedor, J., & Burda, K. (2014). Potential role of carotenoids as antioxidants in human health and disease. *Nutrients*, 6(2), 466–488.
- Bystrická, J., Kavalcová, P., Musilová, J., Vollmannová, A., Tóth, T., Lenková, M. (2015). Carrot (*Daucus carota* L. ssp. *sativus* (Hoffm.) Arcang.) as source of antioxidants. *Acta Agric. Slov.* 105, 303–311.
- Alasalvar, C., Grigor, J. M., Zhang, D., Quantick, P. C., & Shahidi, F. (2001). Comparison of volatiles, phenolics, sugars, antioxidant vitamins, and sensory quality of different colored carrot varieties. *Journal of agricultural and food chemistry*, 49(3), 1410–1416.
- Moldrem, K. L., Li, J., Simon, P. W., & Tanumihardjo, S. A. (2004). Lutein and beta-carotene from lutein-containing yellow carrots are bioavailable in humans. *The American journal of clinical nutrition*, 80(1), 131–136.
- Stange, C., & Rodriguez-Concepcion, M. (2015). Carotenoids in Carrot. In *Pigments in Fruits and Vegetables; Springer Science and Business Media LLC: Berlin/Heidelberg, Germany*, 217–228.
- Hager, T.J., Howard, L.R. (2006). Processing Effects of Carrot Phytonutrients. *Horticultural Science*, 41: 74-79.
- Sharma, K. D., Karki, S., Thakur, N. S., & Attri, S. (2012). Chemical composition, functional properties and processing of carrot-a review. *Journal of food science and technology*, 49(1), 22–32.
- Leja, M., Kamińska, I., Kramer, M., Maksylewicz-Kaul, A., Kammerer, D., Carle, R., & Baranski, R. (2013). The content of phenolic compounds and radical scavenging activity varies with carrot origin and root color. *Plant foods for human nutrition (Dordrecht, Netherlands)*, 68(2), 163–170.

- Ismail, J., Shebaby, W. N., Daher, J., Boulos, J. C., Taleb, R., Daher, C. F., & Mroueh, M. (2023). The Wild Carrot (*Daucus carota*): A Phytochemical and Pharmacological Review. *Plants (Basel, Switzerland)*, 13(1), 93.
- Myojin, C., Enami, N., Nagata, A., Yamaguchi, T., Takamura, H., & Matoba, T. (2008). Changes in the radical-scavenging activity of bitter melon (*Momordica charantia* L.) during freezing and frozen storage with or without blanching. *Journal of food science*, 73(7), C546–C550.
- Phan, M. A. T., Paterson, J., Bucknall, M., & Arcot, J. (2018). Interactions between phytochemicals from fruits and vegetables: Effects on bioactivities and bioavailability. *Critical reviews in food science and nutrition*, 58(8), 1310–1329.
- Zhu, F., Du, B., & Xu, B. (2018). Anti-inflammatory effects of phytochemicals from fruits, vegetables, and food legumes: A review. *Critical reviews in food science and nutrition*, 58(8), 1260–1270.
- Lecerf, J. M., & de Lorgeril, M. (2011). Dietary cholesterol: from physiology to cardiovascular risk. *The British journal of nutrition*, 106(1), 6–14.
- Xu, Z., McClure, S. T., & Appel, L. J. (2018). Dietary Cholesterol Intake and Sources among U.S Adults: Results from National Health and Nutrition Examination Surveys (NHANES), 2001–2014. *Nutrients*, 10(6), 771.
- Wilde, D. W., Massey, K. D., Walker, G. K., Vollmer, A., & Grekin, R. J. (2000). High-fat diet elevates blood pressure and cerebrovascular muscle Ca²⁺ current. *Hypertension (Dallas, Tex.: 1979)*, 35(3), 832–837.
- Ingelsson, E., Schaefer, E. J., Contois, J. H., McNamara, J. R., Sullivan, L., Keyes, M. J., Pencina, M. J., Schoonmaker, C., Wilson, P. W., D'Agostino, R. B., & Vasan, R. S. (2007). Clinical utility of different lipid measures for prediction of coronary heart disease in men and women. *JAMA*, 298(7), 776–785.
- Duan, Y., Zeng, L., Zheng, C., Song, B., Li, F., Kong, X., & Xu, K. (2018). Inflammatory Links Between High Fat Diets and Diseases. *Frontiers in immunology*, 9, 2649.
- Carson, J. A. S., Lichtenstein, A. H., Anderson, C. A. M., Appel, L. J., Kris-Etherton, P. M., Meyer, K. A., Petersen, K., Polonsky, T., Van Horn, L., & American Heart Association Nutrition Committee of the Council on Lifestyle and Cardiometabolic Health; Council on Arteriosclerosis, Thrombosis and Vascular Biology; Council on Cardiovascular and Stroke Nursing; Council on Clinical Cardiology; Council on Peripheral Vascular Disease; and Stroke Council (2020). Dietary Cholesterol and Cardiovascular Risk: A Science Advisory from the American Heart Association. *Circulation*, 141(3), e39–e53.
- Wali, J. A., Jarzebska, N., Raubenheimer, D., Simpson, S. J., Rodionov, R. N., & O'Sullivan, J. F. (2020). Cardio-Metabolic Effects of High-Fat Diets and Their Underlying Mechanisms-A Narrative Review. *Nutrients*, 12(5), 1505.
- Guo, Z., Ali, Q., Abaidullah, M., Gao, Z., Diao, X., Liu, B., Wang, Z., Zhu, X., Cui, Y., Li, D., & Shi, Y. (2022). High fat diet-induced hyperlipidemia and tissue steatosis in rabbits through modulating ileal microbiota. *Applied microbiology and biotechnology*, 106(21), 7187–7207.
- Stellaard F. (2022). From Dietary Cholesterol to Blood Cholesterol, Physiological Lipid Fluxes, and Cholesterol Homeostasis. *Nutrients*, 14(8), 1643.
- Zhao, B., Gan, L., Graubard, B. I., Männistö, S., Albanes, D., & Huang, J. (2022). Associations of Dietary Cholesterol, Serum Cholesterol, and Egg Consumption with Overall and Cause-Specific Mortality: Systematic Review and Updated Meta-Analysis. *Circulation*, 145(20), 1506–1520.
- Kirkpatrick, C. F., Sikand, G., Petersen, K. S., Anderson, C. A. M., Aspary, K. E., Bolick, J. P., Kris-Etherton, P. M., & Maki, K. C. (2023). Nutrition interventions for adults with dyslipidemia: A Clinical Perspective from the National Lipid Association. *Journal of clinical lipidology*, 17(4), 428–451.
- Ayeni, A., Abubakar, A., Aliyu, N., Uhomobhi, L., & Garba, I. (2019). 'Acute and sub-acute toxicity of the crude extracts of the aerial parts of *Daucus carota* L. in laboratory rats'. *Journal of Medicinal Plants for Economic Development*, 3(1), a69.
- Anne Moorhead, S., Welch, R. W., Barbara, M., Livingstone, E., McCourt, M., Burns, A. A., & Dunne, A. (2006). The effects of the fibre content and physical structure of carrots on satiety and subsequent intakes when eaten as part of a mixed meal. *The British journal of nutrition*, 96(3), 587–595.

- Nicolle, C., Cardinault, N., Aprikian, O., Busserolles, J., Grolier, P., Rock, E., Demigné, C., Mazur, A., Scalbert, A., Amouroux, P., & Rémésy, C. (2003). Effect of carrot intake on cholesterol metabolism and on antioxidant status in cholesterol-fed rat. *European journal of nutrition*, 42(5), 254–261.
- Feingold, K. R. (2024). The Effect of Diet on Cardiovascular Disease and Lipid and Lipoprotein Levels. In K. R. Feingold (Eds.) et. al., *Endotext*. MDTText.com, Inc.
- Obia, O., & Eifuobhokhan, J. (2024). Effect of *Justicia carnea* leaf extract on plasma and fecal lipid profile in high-fat diet fed wistar rats. *International Journal of Health and Pharmaceutical Research*. 9(4), 64-70.
- Eifuobhokhan, J., Obia, O., & Charles, C. (2024). *Justicia carnea* leaf extract improves intestinal transit in high-fat diet induced delayed gut motility in wistar rats. *European Journal of Pharmaceutical and Medical Research*. 11(12), 50-54.
- Cheng, H. M., Koutsidis, G., Lodge, J. K., Ashor, A., Siervo, M., & Lara, J. (2017). Tomato and lycopene supplementation and cardiovascular risk factors: A systematic review and meta-analysis. *Atherosclerosis*, 257, 100–108.
- Obia, O., & Asuquo, E.A. (2018). “Endogenous Antioxidant Responses to Dietary Honey Supplementation in Alloxan induced Diabetic Wistar Rats”. *IOSR Journal of Nursing and Health Science (IOSRJNHS)*. 7 (6), 37-40.
- Obia, O., E. Odum, J., & N. Chuemere, A. (2018). Nephroprotective and Antihyperlipedemic Activity of Honey in Alloxan Induced Diabetic Wistar Rats. *International Journal of Biochemistry Research & Review*, 22(1), 1–7.
- Aune D. (2019). Plant Foods, Antioxidant Biomarkers, and the Risk of Cardiovascular Disease, Cancer, and Mortality: A Review of the Evidence. *Advances in nutrition (Bethesda, Md.)*, 10(Suppl_4), S404–S421.
- Chinko, B. C., Pughikumo, D. T., Obia, O., Udeh, W. C., & Hart, V. O. (2023). Honey Attenuates Phenylhydrazine-Induced Hematotoxicity and Oxidative Stress in Male Wistar Rats. *International Blood Research & Reviews*, 14(3), 10–18.
- Marston, N. A., Giugliano, R. P., Im, K., Silverman, M. G., O'Donoghue, M. L., Wiviott, S. D., Ference, B. A., & Sabatine, M. S. (2019). Association Between Triglyceride Lowering and Reduction of Cardiovascular Risk Across Multiple Lipid-Lowering Therapeutic Classes: A Systematic Review and Meta-Regression Analysis of Randomized Controlled Trials. *Circulation*, 140(16), 1308–1317.
- Sirtori, C. R., & Fumagalli, R. (2006). LDL-cholesterol lowering or HDL-cholesterol raising for cardiovascular prevention. A lesson from cholesterol turnover studies and others. *Atherosclerosis*, 186(1), 1–11.
- Kjeldsen, E. W., Nordestgaard, L. T., & Frikke-Schmidt, R. (2021). HDL Cholesterol and Non-Cardiovascular Disease: A Narrative Review. *International journal of molecular sciences*, 22(9), 4547.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/129295>