



Arresting Rancidity in Pearl Millet Flour through Halogen Air Frier Treatment

Vinay G M ^{a*}, S H Akbari ^b, R B Modi ^a, P Srilatha ^a
and Prakasha R ^a

^a Department of Food Processing Technology, College of Food Processing Technology and Bioenergy, Anand Agricultural University, Anand, Gujarat, 388110, India.

^b Department of Food Plant Operation, College of Food Processing Technology and Bioenergy, Anand Agricultural University, Anand, Gujarat, 388110, India.

Authors' contributions

This work was carried out in collaboration among all authors. Author VGM conceptualized the study and did formal analysis, investigation, and performed the methodology. Author SHA supervised the study, reviewed and edited the manuscript, and searched for resources. Author PS did software analysis, data validation, reviewed & edited the manuscript and did data curation. Author RBM supervised the study, searched for resources, reviewed the manuscript and did data validation. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jabb/2025/v28i11858>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/129167>

Original Research Article

Received: 28/10/2024

Accepted: 30/12/2024

Published: 04/01/2025

ABSTRACT

Background: Pearl millet (*Pennisetum glaucum*), a highly nutritious and climate-resilient cereal, is constrained in its utilization due to rapid rancidity and bitterness in its flour. These issues stem from lipid oxidation and the enzymatic activities of lipase and lipoxygenase, which significantly limit the shelf life and consumer acceptability of the flour.

Aim: This study examines the impact of halogen air frying treatment on the enzymatic activity and color characteristics of pearl millet flour.

Study Design: The study was carried out at College of Food Processing Technology and Bio Energy, Anand, in the year 2024.

*Corresponding author: E-mail: vinaygm26@gmail.com;

Methodology: Pearl millet grains were subjected to halogen air fry treatment at temperatures ranging from 110°C to 150°C for durations of 1 to 4 minutes, with a bed thickness of 0.5 cm.

Findings: The results indicate a significant reduction in lipase activity in the treated samples compared to the control, with negligible levels observed at 150°C after 4 minutes. Additionally, lipoxygenase activity was found to be completely inactivated at temperatures of 140°C and 150°C for the same duration. Furthermore, the treated flour demonstrated increased redness and yellowness, highlighting alterations in color values attributable to the heat treatment. These findings suggest that halogen air frying serves as an effective pre-milling treatment for inactivating enzymes responsible for rancidity in pearl millet flour.

Conclusion: This intervention effectively tackles the issue of rancidity while simultaneously prolonging the shelf life of flour, thereby enhancing its overall quality and acceptability. The use of halogen air frying presents a promising and efficient strategy for improving the functional properties of pearl millet flour, facilitating its broader application in food-related industries.

Keywords: Pearl millet; lipase; lipoxygenase; rancidity; shelf life.

1. INTRODUCTION

Pearl millet (*Pennisetum glaucum*), a resilient cereal crop, has been a staple in arid and semi-arid regions for millennia. It has gained significant attention due to its resilience to harsh environmental conditions and high nutritional value. Its origins trace back to the Sahel zone of West Africa. Archaeological evidence indicates its domestication around 2500–2000 BCE in West Africa before introducing to the Indian subcontinent and other parts of Asia (Singh, 2023). Its ability to thrive in regions with high temperatures, low rainfall, and nutrient-poor soils underscores its importance as a climate-resilient crop (Djanaguiraman et al., 2024).

Globally, pearl millet is primarily cultivated in Africa and South Asia. In Africa, countries such as Nigeria, Niger, Mali, and Burkina Faso are leading producers, while in India, states like Rajasthan, Uttar Pradesh, Haryana, Gujarat, and Madhya Pradesh dominate its production. India alone contributes to over 40% of the global pearl millet production by producing 9.62 million tonnes, owing to its adaptability to the semi-arid tropical climate (Directorate of Economics & Statistics, 2022). Its role in ensuring food security for millions in these regions is particularly critical, especially for marginal and resource-poor farmers.

Pearl millet is recognized for its high nutritional profile, which includes high levels of protein, dietary fiber, essential amino acids, and micronutrients like iron and zinc (Vinay and Singh, 2024). Despite these benefits, its consumption faces several barriers, primarily due to issues with the quality and acceptability of processed products. One of the most significant challenges is the rapid onset of rancidity in pearl

millet flour, which severely limits its shelf life (Bhargavi, 2024). Rancidity in pearl millet flour arises from the high lipid content of the grain, particularly in its germ, coupled with the activity of lipolytic and oxidative enzymes such as lipase and lipoxygenase (Ravi and Rana, 2024).

The mechanism of rancidity involves two major pathways: enzymatic and oxidative. Enzymatic rancidity is triggered by lipase, which hydrolyzes triglycerides into free fatty acids. These free fatty acids are further oxidized, leading to the formation of hydroperoxides and secondary volatile compounds like aldehydes and ketones that impart unpleasant odors and flavors (Sruthi and Rao, 2021). Similarly, oxidative rancidity occurs due to the action of lipoxygenase, which catalyzes the oxidation of unsaturated fatty acids, resulting in a deterioration of flavor and quality. The development of bitterness in pearl millet flour is often associated with these processes, compounded by the activity of polyphenol oxidase, which contributes to enzymatic browning and further affects consumer acceptability (Ali et al., 2023).

Processing limitations add to the challenges of utilizing pearl millet in food applications. Traditional methods of milling, such as stone grinding, expose the grain to oxygen and moisture, accelerating rancidity. Moreover, the lack of standardized processing techniques in many production areas limits the development of consistent and high-quality pearl millet products. Efforts to mitigate these challenges have focused on improving processing methods and storage techniques. Thermal treatments, such as hydrothermal, blanching, steaming roasting and extrusion, have been shown to inactivate lipase and lipoxygenase enzymes, thereby enhancing the storage stability of pearl millet flour (Kumar et

al., 2024; Pathare et al., 2024). Similarly, novel techniques such as microwave treatment and controlled-atmosphere packaging have demonstrated potential to delay rancidity and preserve the sensory quality of pearl millet products (Yadav et al., 2012). However, the scalability and economic feasibility of these interventions remain areas of active research.

The present investigation aims to inhibit rancidity-causing enzymes through halogen air fryer treatment with minimal color changes in the pearl millet flour.

2. MATERIALS AND METHODS

The pearl millet of variety Shree Sagar 8484 was procured from the Jibratha Seed Tech LLP, Ahmedabad. The seeds were cleaned to remove immature seeds, stalks, and foreign materials. The standard oleic and linoleic acids were purchased from Sigma Aldrich and the rest of the chemicals were procured from Hi-Media, SRL and Loba Chemie.

2.1 Halogen Air Fry Treatment

The Halogen air fryer (Halogen oven infiniticook 3514I, Usha International Ltd, New Delhi) of 12 L capacity with 1300 W was pre-heated for 1 minute (Fig. 1). The pearl millet grains were spread on the Teflon plate with a bed thickness of 0.5 cm. Later it was placed in a halogen oven and treatment was given at a temperature ranging from 110-150°C for 1-4 minutes. The treated grains were allowed to cool in a desiccator to minimize the moisture gain. After cooling the grains were milled in a laboratory mixer grinder to a fine flour form and sieved in 250 μ size sieve. The flour was packed in metallized laminates (thickness: 110 μ) until further analysis.

2.2 Preparation of Enzyme Extract

The one gram of flour was homogenized with 10 ml of 0.2 M potassium phosphate buffer in a prechilled pestle and mortar. The mixture was centrifuged in a refrigerated centrifuge at 10,000 rpm for 20 min at 4°C. The supernatant was decanted carefully into another test tube and it was used as crude enzyme extract to determine the enzyme activities.

2.3 Lipase Assay

The lipase assay was determined as per the method of Pathare et al. (2024). An aliquot of 100 μ L enzyme extract was combined with 1 mL of

substrate solution consisting of 0.98% (w/v) NaCl, 200 μ L of Tween 20, and an olive oil-water mixture in a 1:3 ratio. The reaction mixture was incubated at 37 °C for 15 minutes, followed by heat inactivation in a water bath at 90 °C for 5 minutes. Subsequently, 5 mL of chloroform was added to the mixture, and it was allowed to incubate at room temperature for 30 minutes. To the lower phase, 2 mL of copper triethanolamine reagent, comprising 1 M triethanolamine, 1 N acetic acid, and 6.45% (w/v) copper nitrate, was thoroughly mixed. This resulted in two distinct color layers: an upper blue layer and a lower yellow layer. Next, 100 μ L of 11 mM diethyl dithiocarbamate (DDC) was added to the lower yellow layer, and its absorbance was measured at 440 nm. A standard curve using oleic acid was prepared, and enzyme activity was determined in terms of mM free fatty acids released per minute per mg of soluble protein.



Fig. 1. Halogen air fryer equipment

2.4 Lipoxxygenase Assay

The assay was performed according to the method mentioned by Goyal and Chugh (2017), using linoleic acid as the substrate. The substrate solution was prepared by gradually adding 0.5 mL of linoleic acid dropwise into 0.5 mL of Tween 20 dissolved in 10 mL of borate buffer at pH 9.0, ensuring thorough mixing to form a fine emulsion. Following this, 1.3 mL of 1 N NaOH was added with continuous stirring to maintain dispersion. Subsequently, 90 mL of borate buffer was mixed into the solution, and the final volume was adjusted to 200 mL with distilled water, yielding a linoleic acid substrate concentration of approximately 7.5 mM.

For the assay, a reaction mixture with a total volume of 2.66 mL was prepared. This mixture

consisted of 2.5 mL of 0.05 M acetate buffer (pH 4.2), 60 μ L of the linoleic acid substrate solution (7.5 mM), and 0.1 mL of enzyme extract. The reaction progress was monitored spectrophotometrically by measuring the increase in absorbance at 234 nm for 2 minutes, using a blank as the reference. The LOX activity was expressed as nanomoles of hydroperoxide produced per minute per milligram of soluble protein (nM HPOD min^{-1} mg^{-1} soluble protein), calculated using the molar extinction coefficient of linoleic acid (25 mM^{-1} cm^{-1}).

2.5 Color Values

The color values of pearl millet flour was measured by Lovibond colorimeter (Model: RT850i) in terms of L^* (Lightness), a^* (redness and greenness) and b^* values (yellowness and blueness). The instrument was calibrated with white and black standards before conducting a test. A flour is filled into a cuvette and placed against the light source (D65). Data were received through the software in terms of L^* (lightness), ranging from 0 (black) to 100 (white), a^* (redness), ranging from +60 (red) to -60 (green), and b^* (yellowness), ranging from +60 (yellow) to -60 (blue) values.

2.6 Statistical Analysis

The results were recorded along with their mean and standard deviation, based on triplicate measurements. One way ANOVA was performed to test the significant difference between the treatments using Tukey (HSD) test.

3. RESULTS AND DISCUSSION

3.1 Effect of Air Fry Treatment on Lipase Activity

The pearl millet flour is susceptible to rancidity by enzymes such as lipase and lipoxygenase and is inactivated at high temperatures. Lipase hydrolyse fat into free fatty acids and hence onset of rancidity occurs. The control pearl millet flour (without treatment) had a lipase activity of 475 mM/min/mg protein. While after giving air fry treatment the lipase activity of the pearl millet flour decreased considerably as depicted in Fig. 2. The lipase activity was decreased to 330

mM/min/mg protein at 110°C for 1 minute, further, its activity decreased to 73.76 mM/min/mg protein after 4 minutes at the same temperature. At 120°C for 1 minute, the lipase activity was found to be 285.43 mM/min/mg protein and it decreased significantly ($p < 0.05$) with time and at 4 minutes the activity was noticed to be 57.89 mM/min/mg protein. The lipase activity of flour treated at 130°C for 1 minute was 226.92 mM/min/mg protein and after 4 minute it was 44.23 mM/min/mg protein. Additionally, the lipase activity of flour at 140°C for 1 minute was 181.78 mM/min/mg protein and decreased to 34.22 mM/min/mg protein after 4 minutes. Furthermore, the lipase activity at 150°C/1 minute was observed to be 173.92 mM/min/mg protein and after 4 minutes, it decreased remarkably to 16.44 mM/min/mg protein. The complete inactivation was not achieved through this treatment but lipase activity was decreased to 16.44 mM/min/mg protein, which is sufficient for the shelf life extension of pearl millet flour for at least 3 months from the 15 days. A similar maximum inhibition was observed by Pathare et al. (2024) in extruded pearl millet flour at 160°C. The enzyme activity decreased with temperature and time due to the denaturation of protein, thereby active site of enzymes was modified and finally its losses its structural integrity.

The free fatty acids formed from hydrolysis of fat by lipase enzyme is further converted into aldehydes and ketones lipoxygenase enzyme. The inactivation of LOX enzyme stabilizes the pearl millet flour by preventing oxidation. The LOX activity of pearl millet flour is depicted in the Fig. 3. As shown in Fig. 2, the control flour had the highest LOX activity of 0.81 nM HPOD/min/mg of protein. As LOX enzyme is more sensitive to heat treatment, all air fry treated flour showed lower enzyme activity (< 0.1 nM HPOD/min/mg of protein). The LOX activity of flour at 110°C/1 minute was 0.09 nM HPOD/min/mg of protein and decreased to 0.02 nM HPOD/min/mg of protein after 4 minutes. Similarly, LOX activity reduced at 120°C/1 minute of 0.08 nM HPOD/min/mg of protein and after 4 minutes decreased to 0.02 nM HPOD/min/mg of protein. The LOX activity was completely not observed at 140°C/4 minutes and 150°C/4 minutes. A similar decrease in the LOX activity was observed by Vinutha et al. (2022), where they have given treatment to flour in NIR lamp (150watt X 4/240 V).




Fig. 2. Lipase activity of pearl millet flour with the air fry treatment. 110, 120, 130, 140 & 150 are the temperature in °C and 1, 2, 3 & 4 are the time in minutes




Fig. 3. Lipoxigenase activity of pearl millet flour

Table 1. Effect of halogen air fry on color values of pearl millet flour

Temperature	Time	L*	a*	b*
Control	-	61.82 ± 0.11 ^b	1.28 ± 0.09 ^{bcd}	12.19 ± 0.06 ^a
110°C	1 minute	67.1 ± 0.19 ^{de}	0.91 ± 0.02 ^a	13.38 ± 0.14 ^c
	2 minutes	69.08 ± 0.74 ^{gh}	2.05 ± 0.06 ^e	16.1 ± 0.05 ^e
	3 minutes	62.03 ± 0.85 ^b	5.86 ± 0.16 ^j	19.35 ± 0.04 ^g
	4 minutes	59.51 ± 0.36 ^a	5.79 ± 0.1 ^j	20.71 ± 0.34 ^{ij}
120°C	1 minute	66.31 ± 0.43 ^{de}	0.94 ± 0.13 ^a	12.57 ± 0.27 ^{ab}
	2 minutes	69.21 ± 0.49 ^{gh}	1.32 ± 0.08 ^{cd}	14.43 ± 0.26 ^d
	3 minutes	66.47 ± 0.26 ^{de}	3.98 ± 0.07 ^g	19.39 ± 0.13 ^g
	4 minutes	61.59 ± 0.14 ^b	6.3 ± 0.1 ^k	21 ± 0.25 ^{ij}
130°C	1 minute	68.87 ± 0.06 ^{gh}	1.07 ± 0.04 ^{abc}	13.43 ± 0.12 ^c
	2 minutes	69.94 ± 0.1 ^h	1.43 ± 0.07 ^d	14.94 ± 0.03 ^d
	3 minutes	66.59 ± 0.02 ^{de}	4.72 ± 0.02 ^h	20.57 ± 0.03 ^{hi}
	4 minutes	63.91 ± 0.17 ^c	5.15 ± 0.02 ⁱ	20.34 ± 0.06 ^{ij}
140°C	1 minute	68.31 ± 0.23 ^{fg}	0.98 ± 0 ^a	13.22 ± 0.18 ^c
	2 minutes	69.79 ± 0.52 ^h	1.31 ± 0.07 ^{cd}	14.73 ± 0.29 ^d
	3 minutes	67.41 ± 0.38 ^{ef}	4.15 ± 0.07 ^g	19.83 ± 0.24 ^{gh}
	4 minutes	62.2 ± 0.18 ^b	6.04 ± 0.01 ^{jk}	20.9 ± 0.02 ^{ij}
150°C	1 minute	66.24 ± 0.52 ^d	1.02 ± 0.09 ^{ab}	12.37 ± 0.35 ^{ab}
	2 minutes	67.26 ± 0.22 ^{def}	1.13 ± 0.01	13 ± 0.04 ^{bc}
	3 minutes	68.81 ± 0.13 ^{gh}	3.02 ± 0.18 ^f	18.05 ± 0.41 ^f
	4 minutes	63.81 ± 0.01 ^c	5.41 ± 0.03 ⁱ	20.85 ± 0.03 ^{ij}

Data are represented as mean ± SD (n=3). The different superscript of letters in a column significantly varies (p<0.05)

3.2 Effect on Color Values

The color values indicate the changes in the appearance of pearl millet flour after undergoing halogen air fry treatment. These values, as presented in Table 1, provide insights into the impact of processing on the flour's visual characteristics. The lightness value (L*) of the untreated control sample was 61.82, which significantly decreased after treatment. However, the variations in lightness did not follow a consistent trend across different treatment conditions. The positive a* values indicate the redness of the samples; the control sample had an a* value of 1.28, while the redness of all treated samples increased significantly (p < 0.05) with higher temperatures and longer treatment times. Similarly, the b* values, which represent the yellowness of the flour, showed a notable increase. The untreated control had a b* value of 12.19, and this value significantly (p < 0.05) increased with treatment time and temperature. These changes in redness and yellowness suggest that higher temperatures induce Maillard reactions or caramelization, leading to darker, more intense color changes in the flour. Overall,

the results demonstrate that high-temperature treatment not only alters the enzyme activity but also significantly impacts the visual properties of the flour, which may influence consumer perception and product quality.

4. CONCLUSION

Pearl millet utilization is limited by the rapid onset of rancidity and bitterness in its flour due to the enzymatic activities of lipase and lipoxygenase. This study demonstrated that halogen air fry treatment effectively addresses these issues by significantly reducing lipase activity and completely inactivating lipoxygenase at 150°C for 4 minutes. The treatment also altered the flour's color properties, increasing redness and yellowness, which may enhance its visual appeal. These findings suggest that halogen air frying is a practical pre-milling treatment for extending the shelf life and improving the sensory quality of pearl millet flour. By inactivating rancidity-causing enzymes, this method provides a simple and efficient solution for overcoming a key barrier to the broader utilization of pearl millet. The approach holds

promise for enhancing the functional and economic value of pearl millet, paving the way for its greater application in food systems while maintaining its nutritional benefits. Future studies could focus on optimizing processing conditions with the moisture content can enhance quality and consumer acceptability.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology.

Details of the AI usage are given below:

1. ChatGPT

DATA AVAILABILITY

The data obtained during the study are available from the corresponding author upon reasonable request.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Bhargavi, H. A., Singh, S. P., Goswami, S., Yadav, S., Aavula, N., Shashikumara, P., ... & Singh, N. (2024). Deciphering the genetic variability for biochemical parameters influencing rancidity of *pearl millet* (*Pennisetum glaucum* L. R. Br.) flour in a set of highly diverse lines and their categorization using rancidity matrix. *Journal of Food Composition and Analysis*, 128, 106035. <https://doi.org/10.1016/j.jfca.2024.106035>
- Directorate of Economics & Statistics. (2022). *Agriculture statistics at a glance*. Ministry of Agricultural and Family Welfare, Government of India, 36-37.
- Djanaguiraman, M., Priyanka, A. S., Vaishnavi, S. J., Perumal, R., Ciampitti, I. A., & Prasad, P. V. (2024). Impact of drought and high-temperature stresses on growth and development stages, physiological, reproductive, and yield traits on *pearl millet*. In *Pearl Millet: A Resilient Cereal Crop for Food, Nutrition, and Climate Security* (pp. 249-276).
- Goyal, P., & Chugh, L. K. (2017). Shelf life determinants and enzyme activities of *pearl millet*: A comparison of changes in stored flour of hybrids, CMS lines, inbreds, and composites. *Journal of Food Science and Technology*, 54, 3161-3169. <https://doi.org/10.1007/s13197-017-2760-6>
- Kumar, R. R., Singh, N., Goswami, S., Vinutha, T., Singh, S. P., Mishra, G. P., ... & Tyagi, A. (2024). Hydrothermal infra-red (HT-IR): The most effective technology for enhancing the shelf-life of *pearl millet* flour without compromising the nutrient density and flour quality. *Journal of Plant Biochemistry and Biotechnology*, 1-15. <https://doi.org/10.1007/s13562-023-00753-8>
- Pathare, A. M., Singhal, R. S., Rao, B. D., & Gokhale, J. S. (2024). Extrusion inhibits lipolytic enzymes in *pearl millet* (*Pennisetum glaucum*) flour. *Innovative Food Science & Emerging Technologies*, 93, 103605. <https://doi.org/10.1016/j.ifset.2023.103605>
- Ravi, J. L., & Rana, S. S. (2024). Maximizing the nutritional benefits and prolonging the shelf life of millets through effective processing techniques: A review. *ACS Omega*, 9(37), 38327-38347. <https://doi.org/10.1021/acsomega.3c05978>
- Singh, A. K. (2023). Early presence/introduction of African and East Asian millets in India: Integral to traditional agriculture. *The Nucleus*, 66(3), 261-271. <https://doi.org/10.1007/s13237-023-00425-9>
- Sruthi, N. U., & Rao, P. S. (2021). Effect of processing on storage stability of millet flour: A review. *Trends in Food Science & Technology*, 112, 58-74. <https://doi.org/10.1016/j.tifs.2021.03.041>
- Vinay, G. M., & Singh, A. K. (2024). Effect of ghee residue powder and *pearl millet* flour substitution on rheological, textural, and sensorial characteristics of eggless muffin. *Journal of Food Processing and Preservation*, 2024(1), 5519265. <https://doi.org/10.1111/jfpp.5519265>
- Vinutha, T., Kumar, D., Bansal, N., Krishnan, V., Goswami, S., Kumar, R. R., ... & Praveen, S. (2022). Thermal treatments reduce rancidity and modulate structural and

digestive properties of starch in *pearl millet* flour. *International Journal of Biological Macromolecules*, 195, 207-216. <https://doi.org/10.1016/j.ijbiomac.2021.12.176>

Yadav, D. N., Anand, T., Kaur, J., & Singh, A. K. (2012). Improved storage stability of *pearl millet* flour through microwave treatment. *Agricultural Research*, 1, 399-404. <https://doi.org/10.1007/s40003-012-0038-x>

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/129167>